

quatrième série - tome 56 fascicule 4 juillet-août 2023

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Olivier BENOIST & Olivier WITTENBERG

Intermediate Jacobians and rationality over arbitrary fields

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

YVES DE CORNULIER

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE
de 1883 à 1888 par H. DEBRAY
de 1889 à 1900 par C. HERMITE
de 1901 à 1917 par G. DARBOUX
de 1918 à 1941 par É. PICARD
de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} mai 2023

S. CANTAT G. GIACOMIN
G. CARRON D. HÄFNER
Y. CORNULIER D. HARARI
F. DÉGLISE C. IMBERT
A. DUCROS S. MOREL
B. FAYAD P. SHAN
J. FRESÁN

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.
Email : annaes@ens.fr

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France
Case 916 - Luminy
13288 Marseille Cedex 09
Tél. : (33) 04 91 26 74 64. Fax : (33) 04 91 41 17 51
Email : abonnements@smf.emath.fr

Tarifs

Abonnement électronique : 459 euros.
Abonnement avec supplément papier :
Europe : 646 €. Hors Europe : 730 € (\$ 985). Vente au numéro : 77 €.

© 2023 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Fabien Durand
Périodicité : 6 n^{os} / an

INTERMEDIATE JACOBIANS AND RATIONALITY OVER ARBITRARY FIELDS

BY OLIVIER BENOIST AND OLIVIER WITTENBERG

ABSTRACT. – We prove that a three-dimensional smooth complete intersection of two quadrics over a field k is k -rational if and only if it contains a line defined over k . To do so, we develop a theory of intermediate Jacobians for geometrically rational threefolds over arbitrary, not necessarily perfect, fields. As a consequence, we obtain the first examples of smooth projective varieties over a field k which have a k -point, and are rational over a purely inseparable field extension of k , but not over k .

RÉSUMÉ. – Nous démontrons qu'une intersection complète lisse de deux quadriques de dimension 3 sur un corps k est k -rationnelle si et seulement si elle contient une droite définie sur k . À cet effet, nous développons une théorie des jacobiniennes intermédiaires pour les variétés géométriquement rationnelles de dimension 3 sur des corps quelconques, non nécessairement parfaits. Comme conséquence, nous obtenons les premiers exemples de variétés projectives lisses sur un corps k qui ont un k -point, et sont rationnelles sur une extension de corps purement inséparable de k , mais pas sur k .

Introduction

Let k be a field. A variety X of dimension n over k is said to be k -rational (resp. k -unirational, resp. separably k -unirational) if there exists a birational map (resp. a dominant rational map, resp. a dominant and separable rational map) $\mathbf{A}_k^n \dashrightarrow X$.

This article is devoted to studying the k -rationality of threefolds over k . Our main result answers positively a conjecture of Kuznetsov and Prokhorov.

THEOREM A (Theorem 4.7). – *Let $X \subset \mathbf{P}_k^5$ be a smooth complete intersection of two quadrics. Then X is k -rational if and only if it contains a line defined over k .*

The question of the validity of Theorem A goes back to Auel, Bernardara and Bolognesi [5, Question 5.3.2 (3)], who raised it when k is a rational function field in one variable over an algebraically closed field.

Using the fact that varieties X as in Theorem A are separably k -unirational if and only if they have a k -point (see Theorem 4.8), we obtain new counterexamples to the Lüroth problem over non-closed fields.

THEOREM B (Theorem 4.14). – *For any algebraically closed field κ , there exists a three-dimensional smooth complete intersection of two quadrics $X \subset \mathbf{P}_{\kappa((t))}^5$ which is separably $\kappa((t))$ -unirational, $\kappa((t^{\frac{1}{2}}))$ -rational, but not $\kappa((t))$ -rational.*

When κ has characteristic 2, Theorem B yields the first examples of smooth projective varieties over a field k which have a k -point and are rational over the perfect closure of k , but which are not k -rational (see Remarks 4.15 (iii) and (iv)).

Theorem A may be compared to the classical fact that a smooth quadric over k is k -rational if and only if it has a k -point. However, although it is easy to check that a smooth projective k -rational variety has a k -point, the fact that a k -rational three-dimensional smooth complete intersection of two quadrics X necessarily contains a k -line is highly non-trivial. To prove it, we rely on obstructions to the k -rationality of X arising from a study of its intermediate Jacobian.

Such obstructions go back to the seminal work of Clemens and Griffiths [15]: if a smooth projective threefold over \mathbf{C} is \mathbf{C} -rational, then its intermediate Jacobian is isomorphic, as a principally polarized abelian variety over \mathbf{C} , to the Jacobian of a (not necessarily connected) smooth projective curve. This implication was used in [15] to show that smooth cubic threefolds over \mathbf{C} are never \mathbf{C} -rational, and was later applied to show the irrationality of several other classes of complex threefolds (see for instance [6]). The work of Clemens and Griffiths was extended by Murre [71] to algebraically closed fields of any characteristic different from 2.

More recently, we implemented the arguments of Clemens and Griffiths over arbitrary perfect fields k [7]. By exploiting the fact that the intermediate Jacobian may be isomorphic to the Jacobian of a smooth projective curve over \bar{k} while not being so over k , we produced new examples of varieties over k that are \bar{k} -rational but not k -rational.

Hassett and Tschinkel [46] subsequently noticed that over a non-closed field k , the intermediate Jacobian carries further obstructions to k -rationality: if X is a smooth projective k -rational threefold, then not only is its intermediate Jacobian isomorphic to the Jacobian $\mathbf{Pic}^0(C)$ of a smooth projective curve C over k , but in addition, assuming for simplicity that C is geometrically connected of genus ≥ 2 , the $\mathbf{Pic}^0(C)$ -torsors associated with $\mathrm{Aut}(\bar{k}/k)$ -invariant algebraic equivalence classes of codimension 2 cycles on X are of the form $\mathbf{Pic}^\alpha(C)$ for some $\alpha \in \mathbf{Z}$. When X is a smooth three-dimensional complete intersection of two quadrics, they used these obstructions in combination with the natural identification of the variety of lines of X with a torsor under the intermediate Jacobian of X , and with the work of Wang [82], to prove Theorem A when $k = \mathbf{R}$ [46, Theorem 36] (and later [45] over subfields of \mathbf{C}).

The aim of the present article is to extend these arguments to arbitrary fields.

Applications to k -rationality criteria for other classes of \bar{k} -rational threefolds appear in the work of Kuznetsov and Prokhorov [61].

So far, we have been imprecise about what we call the intermediate Jacobian of a smooth projective threefold X over k .

If $k = \mathbf{C}$, one can use Griffiths' intermediate Jacobian $J^3 X$ constructed by transcendental means. This is the original path taken by Clemens and Griffiths [15]. The algebraic part of Griffiths' intermediate Jacobian has been shown to descend to subfields $k \subseteq \mathbf{C}$ by Achter, Casalaina-Martin and Vial [3, Theorem B]; the resulting k -structure on $J^3 X$ is the one used in [45].

Over algebraically closed fields k of arbitrary characteristic, a different construction of an intermediate Jacobian $\text{Ab}^2(X)$, based on codimension 2 cycles, was provided by Murre [72, Theorem A p. 226] (see also [54]). This cycle-theoretic approach to intermediate Jacobians had already been applied by him to rationality problems (see [71]).

Over a perfect field k , the universal property satisfied by Murre's intermediate Jacobian $\text{Ab}^2(X_{\bar{k}})$ induces a Galois descent datum on $\text{Ab}^2(X_{\bar{k}})$, thus yielding a k -form $\text{Ab}^2(X)$ of $\text{Ab}^2(X_{\bar{k}})$ [3, Theorem 4.4]. It is this intermediate Jacobian $\text{Ab}^2(X)$, which coincides with $J^3 X$ when $k \subseteq \mathbf{C}$, that we used in [7].

Over an imperfect field k , one runs into the difficulty that Murre's definition of $\text{Ab}^2(X_{\bar{k}})$ does not give rise to a \bar{k}/k -descent datum on $\text{Ab}^2(X_{\bar{k}})$. Achter, Casalaina-Martin and Vial still prove, in [4], the existence of an algebraic representative $\text{Ab}^2(X)$ for algebraically trivial codimension 2 cycles on X (see §1.2 of *op. cit.* for the definition). However, when k is imperfect, it is not known whether $\text{Ab}^2(X)_{\bar{k}}$ is isomorphic to $\text{Ab}^2(X_{\bar{k}})$. For this reason, we do not know how to construct on $\text{Ab}^2(X)$ the principal polarization that is so crucial to the Clemens-Griffiths method.

To overcome this difficulty and prove Theorem A in full generality, we provide, over an arbitrary field k , an entirely new construction of an intermediate Jacobian.

Our point of view is inspired by Grothendieck's definition of the Picard scheme (for which see [33], [12, Chapter 8], [56]). With any smooth projective \bar{k} -rational threefold X over k , we associate a functor $\text{CH}_{X/k, \text{fppf}}^2 : (\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$ endowed with a natural bijection $\text{CH}^2(X_{\bar{k}}) \xrightarrow{\sim} \text{CH}_{X/k, \text{fppf}}^2(\bar{k})$ (see Definition 2.9 and (3.1)). The functor $\text{CH}_{X/k, \text{fppf}}^2$ is an analogue, for codimension 2 cycles, of the Picard functor $\text{Pic}_{X/k, \text{fppf}}$.

Too naive attempts to define the functor $\text{CH}_{X/k, \text{fppf}}^2$ on the category of k -schemes, such as the formula " $T \mapsto \text{CH}^2(X_T)$ ", fail as Chow groups of possibly singular schemes are not even contravariant with respect to arbitrary morphisms: one would need to use a contravariant variant of Chow groups (see Remark 3.2 (ii)). To solve this issue, we view Chow groups of codimension ≤ 2 as subquotients of K -theory by means of the Chern character, and we define $\text{CH}_{X/k, \text{fppf}}^2$ as an appropriate subquotient of (the fppf sheafification of) the functor $T \mapsto \mathbf{K}_0(X_T)$. That this procedure gives rise to the correct functor, even integrally, is a consequence of the Riemann-Roch theorem without denominators [53].

We show that $\text{CH}_{X/k, \text{fppf}}^2$ is represented by a smooth k -group scheme $\mathbf{CH}_{X/k}^2$ (Theorem 3.1 (i)). Our functorial approach is crucial for this, as it allows us to argue by fppf descent from a possibly inseparable finite extension l of k such that X is l -rational. By construction, there is a natural isomorphism $\mathbf{CH}_{X/l}^2 \simeq (\mathbf{CH}_{X/k}^2)_l$ for all field extensions l of k .

The k -group scheme that we use as a substitute for the intermediate Jacobian of X is then the identity component $(\mathbf{CH}_{X/k}^2)^0$ of $\mathbf{CH}_{X/k}^2$, which is an abelian variety (Theorem 3.1 (ii)). We hope that this functorial perspective on intermediate Jacobians may have other applications (to intermediate Jacobians in families, to deformations of algebraic cycles).

Establishing an identification $(\mathbf{CH}_{X/k}^2)^0_{\bar{k}} \simeq \mathbf{Ab}^2(X_{\bar{k}})$ (Theorem 3.1 (vi)) and using the principal polarization on $\mathbf{Ab}^2(X_{\bar{k}})$ constructed in [7], we endow $(\mathbf{CH}_{X/k}^2)^0$ with a canonical principal polarization, which paves the way for applications to rationality questions. Let us now state the most general obstruction to the k -rationality of a smooth projective threefold that we obtain by analyzing $\mathbf{CH}_{X/k}^2$.

THEOREM C (Theorem 3.1 (vii)). – *Let X be a smooth projective k -rational threefold over k . Then there exists a smooth projective curve B over k such that the k -group scheme $\mathbf{CH}_{X/k}^2$ can be realized as a direct factor of $\mathbf{Pic}_{B/k}$ in a way that respects the canonical principal polarizations.*

In Theorem 3.11, we deduce from Theorem C more concrete obstructions to the k -rationality of X , pertaining to the Néron-Severi group $\mathbf{NS}^2(X_{\bar{k}})$ of algebraic equivalence classes of codimension 2 cycles on $X_{\bar{k}}$, to the principally polarized abelian variety $(\mathbf{CH}_{X/k}^2)^0$, and to the $(\mathbf{CH}_{X/k}^2)^0$ -torsors that are of the form $(\mathbf{CH}_{X/k}^2)^\alpha$ for some $\alpha \in (\mathbf{CH}_{X/k}^2 / (\mathbf{CH}_{X/k}^2)^0)(k) = \mathbf{NS}^2(X_{\bar{k}})^{\mathbf{Aut}(\bar{k}/k)}$.

The principle of the proof of Theorem C goes back to Clemens and Griffiths. Since X is k -rational, it can be obtained from \mathbf{P}_k^3 by a composition of blow-ups of regular curves and of closed points, followed by a contraction. The curve B whose existence is predicted by Theorem C is roughly the union of the blown-up curves. This works perfectly well if k is perfect. If k is imperfect, however, some of the blown-up curves may be regular but not smooth over k . It is nevertheless very important, in view of the application to Theorem A, that the curve B appearing in the statement of Theorem C be smooth over k . To prove Theorem C as stated, one thus has to show that the contributions of these non-smooth regular curves are canceled out by the final contraction. This non-trivial fact relies on a complete understanding of which Jacobians of proper reduced curves over k split as the product of an affine group scheme and of an abelian variety over k (Theorem 1.7).

The text is organized as follows. Sections 1 and 2 gather preliminaries, concerning respectively group schemes and K -theory. Section 2 contains in particular the definition of the above-mentioned functor $\mathbf{CH}_{X/k, \text{fppf}}^2$ associated with a smooth projective \bar{k} -rational threefold X over k (Definition 2.9). In Section 3, we prove that this functor is representable and study the k -group scheme $\mathbf{CH}_{X/k}^2$ that represents it (Theorem 3.1). A number of obstructions to the k -rationality of X are then derived (Theorem 3.1 (vii) and Theorem 3.11). Section 4 is devoted to applications to three-dimensional smooth complete intersections of two quadrics: we compute $\mathbf{CH}_{X/k}^2$ entirely (Theorem 4.5), deduce the irrationality criterion that is our main theorem (Theorem 4.7), and apply the criterion to examples over Laurent series fields (Theorem 4.14).

Acknowledgements

Over a perfect field, the results of this article were obtained during the “Rationality of Algebraic Varieties” conference held on Schiermonnikoog Island in April 2019 after reading the arguments of Hassett and Tschinkel [46] over the reals. We are grateful to Jean-Louis Colliot-Thélène and to Sasha Kuznetsov for encouraging us to write them up. Finally, we thank Asher Auel, Marcello Bernardara and Michele Bolognesi for their comments and their interest, and the referees for their work and their helpful suggestions.

Notation and conventions

We fix a field k and an algebraic closure \bar{k} of k . Let k_p be the perfect closure of k in \bar{k} and $\Gamma_k := \text{Gal}(\bar{k}/k_p) = \text{Aut}(\bar{k}/k)$ be the absolute Galois group of k_p . If X and T are two k -schemes, we set $X_T := X \times_k T$. A *variety* over k is a separated scheme of finite type over k ; a *curve* is a variety of pure dimension 1. If G is a group scheme locally of finite type over k , we denote by G^0 its identity component. If X is a smooth proper variety over k , we let $\text{CH}^c(X_{\bar{k}})_{\text{alg}} \subseteq \text{CH}^c(X_{\bar{k}})$ be the subgroup of algebraically trivial codimension c cycle classes and define $\text{NS}^c(X_{\bar{k}}) := \text{CH}^c(X_{\bar{k}}) / \text{CH}^c(X_{\bar{k}})_{\text{alg}}$.

We use *qcqs* as a shorthand for quasi-compact and quasi-separated. We denote by (Sch/k) the category of qcqs k -schemes and by (Ab) the category of abelian groups. If X is a commutative k -group scheme, we let $\Phi_X : (\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$ be the functor given by $\Phi_X(T) = \text{Hom}_k(T, X)$. The functor $\Phi : X \mapsto \Phi_X$, from the category of commutative k -group schemes to the category of functors $(\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$, is fully faithful, by Yoneda’s lemma and since all schemes are covered by affine (hence qcqs) open subschemes. We say that a functor $(\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$ is *representable* if it is isomorphic to Φ_X for some commutative k -group scheme X , which need not be qcqs. The functor $\mathbf{Z} : (\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$ sending $T \in (\text{Sch}/k)$ to the group $\mathbf{Z}(T)$ of locally constant maps $T \rightarrow \mathbf{Z}$ is represented by the constant k -group scheme \mathbf{Z} .

We refer to [38, VII, Définition 1.4] for the definition of a *regular immersion* of schemes that we use, based on the Koszul complex. A closed immersion of regular schemes is always regular. A morphism of schemes $f : X \rightarrow Y$ is said to be a *local complete intersection* or *lci* if it can be factored, locally on X , as the composition of a regular immersion and of a smooth morphism [38, VIII, Définition 1.1].

If ℓ is a prime number and M is a \mathbf{Z} -module, we let $M\{\ell\} \subseteq M$ be the ℓ -primary torsion subgroup of M , and $T_\ell(M) := \text{Hom}(\mathbf{Q}_\ell/\mathbf{Z}_\ell, M)$ and $V_\ell(M) := T_\ell(M)[1/\ell]$ be the ℓ -adic Tate modules of M . If G is a commutative k -group scheme, we set $T_\ell G := T_\ell(G(\bar{k}))$ and $V_\ell G := V_\ell(G(\bar{k}))$.

1. Group schemes

We first collect miscellaneous information concerning group schemes. The main new result of this section is Theorem 1.7.

1.1. Chevalley's theorem

If G is a connected smooth group scheme over k , there is a unique short exact sequence

$$(1.1) \quad 0 \rightarrow L(G_{k_p}) \rightarrow G_{k_p} \rightarrow A(G_{k_p}) \rightarrow 0$$

of group schemes over k_p , where $L(G_{k_p})$ is smooth, connected and affine and where $A(G_{k_p})$ is an abelian variety. This statement was proved by Chevalley [14], see [18, Theorem 1.1] for a modern proof.

1.2. Principal polarizations

Recall that a *principal polarization* on an abelian variety A over k_p is an ample class $\theta \in \text{NS}(A_{\bar{k}})^{\Gamma_k}$ whose associated isogeny $A_{\bar{k}} \rightarrow \hat{A}_{\bar{k}}$ is an isomorphism, that a principally polarized abelian variety A over k_p is a product of indecomposable principally polarized abelian varieties over k_p and that the factors of this decomposition are unique as subvarieties of A (see [7, §2.1]).

We define a *polarization* (resp. a *principal polarization*) of a smooth commutative group scheme G over k to be a polarization (resp. a principal polarization) of the abelian variety $A(G_{k_p}^0)$ over k_p . If G is endowed with a polarization θ , we say that a smooth subgroup scheme $H \subseteq G$ is a *polarized direct factor* of G if there exists a subgroup scheme $H' \subseteq G$ such that the canonical morphism $\iota : H \times H' \xrightarrow{\sim} G$ is an isomorphism and such that $\iota^*\theta = \pi^*\theta|_{A(H_{k_p}^0)} + \pi'^*\theta|_{A(H'_{k_p}^0)}$, where π and π' denote the projections of $A(H_{k_p}^0) \times A(H'_{k_p}^0)$ onto $A(H_{k_p}^0)$ and $A(H'_{k_p}^0)$. If in addition θ is a principal polarization, then so are $\pi^*\theta|_{A(H_{k_p}^0)}$ and $\pi'^*\theta|_{A(H'_{k_p}^0)}$; in this case, we also speak of a *principally polarized direct factor*.

1.3. Locally constant functions

For a variety X over k , consider the functor

$$\begin{aligned} \mathbf{Z}_{X/k} : (\text{Sch}/k)^{\text{op}} &\rightarrow (\text{Ab}) \\ T &\mapsto \mathbf{Z}(X_T). \end{aligned}$$

Equivalently $\mathbf{Z}_{X/k}$ is the push-forward of the constant sheaf \mathbf{Z} by the structural morphism $X \rightarrow \text{Spec}(k)$ (and hence is an fpqc sheaf). Let $\pi_0(X/k)$ denote the étale k -scheme of connected components of X , defined in [26, §I.4, Définition 6.6]. The Weil restriction of scalars $\text{Res}_{\pi_0(X/k)/k}\mathbf{Z}$ exists as a k -scheme by [12, 7.6/4].

PROPOSITION 1.1. – *Let X be a variety over k . Then the functor $\mathbf{Z}_{X/k}$ is canonically represented by the Weil restriction of scalars $\text{Res}_{\pi_0(X/k)/k}\mathbf{Z}$.*

Proof. – Recall that there is a canonical faithfully flat morphism $q_X : X \rightarrow \pi_0(X/k)$ whose fibers are geometrically connected [26, §I.4, Propositions 6.5 and 6.7]. For any $T \in (\text{Sch}/k)$, the resulting morphism $X_T \rightarrow \pi_0(X/k)_T$ is surjective, has connected fibers, and is open [35, Théorème 2.4.6]; therefore q_X induces a bijection between the sets of connected components of X_T and of $\pi_0(X/k)_T$. As a consequence, the homomorphism $\mathbf{Z}(\pi_0(X/k)_T) \rightarrow \mathbf{Z}(X_T)$ is an isomorphism, and hence so is the morphism of functors $\text{Res}_{\pi_0(X/k)/k}\mathbf{Z} \rightarrow \mathbf{Z}_{X/k}$ that it induces when T varies. \square

REMARKS 1.2. – (i) We will usually denote by $\mathbf{Z}_{X/k}$, rather than by $\text{Res}_{\pi_0(X/k)/k}\mathbf{Z}$, the group scheme that $\mathbf{Z}_{X/k}$ represents.

(ii) Proposition 1.1 shows that a morphism $p : X' \rightarrow X$ of varieties over k that induces a bijection between the sets of connected components of $X_{\bar{k}}$ and of $X'_{\bar{k}}$ gives rise to an isomorphism $\mathbf{Z}_{X/k} \xrightarrow{\sim} \mathbf{Z}_{X'/k}$.

1.4. Picard schemes

The *absolute Picard functor* of a proper variety X over k is

$$\begin{aligned} \text{Pic}_{X/k} : (\text{Sch}/k)^{\text{op}} &\rightarrow (\text{Ab}) \\ T &\mapsto \text{Pic}(X_T). \end{aligned}$$

Beware that our notation differs slightly from that of [56, §9.2].

If $\tau \in \{\text{Zar}, \acute{\text{e}}\text{t}, \text{fppf}\}$, we denote the sheafification of $\text{Pic}_{X/k}$ for the corresponding (Zariski, étale or fppf) topology by $\text{Pic}_{X/k, \tau}$. The functors $\text{Pic}_{X/k, \acute{\text{e}}\text{t}}$ and $\text{Pic}_{X/k, \text{fppf}}$ are equal [12, 8.1 p. 203] and are represented by a group scheme locally of finite type over k [12, 8.2/3] which we denote by $\mathbf{Pic}_{X/k}$: the *Picard scheme* of X . These two functors contain $\text{Pic}_{X/k, \text{Zar}}$ and $T \mapsto \text{Pic}(X_T)/\text{Pic}(T)$ as subfunctors if $H^0(X, \mathcal{O}_X) = k$, and they coincide with them if in addition $X(k) \neq \emptyset$ (see [56, Theorem 9.2.5] and [34, Proposition 7.8.6]), for instance if X is connected and reduced and $k = \bar{k}$.

1.4.1. *Picard schemes of blow-ups.* – In §1.4.1, we consider the following situation. We fix a regular closed immersion $i : Y \rightarrow X$ of qcqs schemes of pure codimension $c \geq 2$, we let $p : X' \rightarrow X$ be the blow-up of X along Y with exceptional divisor Y' , and we let $p' : Y' \rightarrow Y$ and $i' : Y' \rightarrow X'$ be the natural morphisms. The morphism $p' : Y' \rightarrow Y$ is a projective bundle of relative dimension $c - 1 \geq 1$ (see [79, §1.2]). In this setting, we study the group morphism

$$(1.2) \quad \begin{aligned} \text{Pic}(X) \times \mathbf{Z}(Y) &\rightarrow \text{Pic}(X') \\ (\mathcal{L}, \psi) &\mapsto p^* \mathcal{L} \otimes_{\mathcal{O}_{X'}} \left(- \sum_{n \in \mathbf{Z}} n [p^{-1}(\psi^{-1}(n))] \right). \end{aligned}$$

PROPOSITION 1.3. – *Under the hypotheses of §1.4.1, the map (1.2) is bijective.*

Proof. – By absolute noetherian approximation [80, Theorem C.9] and the limit arguments of [36, §8], we may assume that X is noetherian. If $\mathcal{N} \in \text{Pic}(X')$, the function $\psi : Y \rightarrow \mathbf{Z}$ such that $\mathcal{N}|_{X'_y} \simeq \mathcal{O}_{X'_y}(\psi(y))$ for all $y \in Y$ is locally constant. (Indeed, for $n \gg 0$, the \mathcal{O}_Y -module $p'_*(\mathcal{N}|_{Y'})_n$ is locally free and its formation commutes with base change, by [44, III, Theorems 8.8 and 12.11].) Since $\mathcal{O}_{X'}(-Y')|_{X'_y} \simeq \mathcal{O}_{X'_y}(1)$ for all $y \in Y$ (see [79, §1.2]), it follows that $\mathcal{N} \otimes_{\mathcal{O}_{X'}} (\sum_{n \in \mathbf{Z}} n [p^{-1}(\psi^{-1}(n))])$ is trivial on the fibers of p , and that ψ is the unique function with this property. It remains to show that $p^* : \text{Pic}(X) \rightarrow \text{Pic}(X')$ is injective with image the subgroup of isomorphism classes of line bundles that are trivial on the fibers of p . The injectivity follows from [79, Lemme 2.3 (a)], and the description of the image from Lemma 1.4 (iii) below. \square

LEMMA 1.4. – *Under the hypotheses of §1.4.1, assume that X is noetherian and let \mathcal{N} be a line bundle on X' such that $\mathcal{N}|_{X'_y} \simeq \mathcal{O}_{X'_y}$ for all $y \in Y$. For any integer n , set $\mathcal{N}(n) = \mathcal{N} \otimes \mathcal{O}_{X'}(-nY')$. Then:*

- (i) *For all $j \geq 1$ and $n \geq 0$, the sheaf $R^j p_*(\mathcal{N}(n))$ vanishes.*
- (ii) *For all $n \geq 0$, the natural morphism $p^* p_*(\mathcal{N}(n)) \rightarrow \mathcal{N}(n)$ is surjective.*
- (iii) *The sheaf $p_* \mathcal{N}$ is invertible and $p^* p_* \mathcal{N} \rightarrow \mathcal{N}$ is an isomorphism.*

Proof. – By [44, III, Theorem 8.8 (c)], assertion (i) holds for $n \gg 0$. To prove (i) for all $n \geq 0$ by descending induction on n , we consider for $j \geq 1$ the exact sequence

$$R^j p_* \mathcal{N}(n) \rightarrow R^j p_* \mathcal{N}(n-1) \rightarrow R^j p_*(\mathcal{N}(n-1)|_{Y'})$$

and note that $R^j p'_*(\mathcal{N}(n-1)|_{Y'}) = 0$ for $n \geq 1$ by cohomology and base change [44, III, Theorem 12.11].

Assertion (ii) holds for all $n \gg 0$ by [44, III, Theorem 8.8 (a)]. To prove (ii) for all $n \geq 0$ by descending induction on n , we consider the natural commutative diagram with exact rows

$$\begin{array}{ccccccc} p^* p_* \mathcal{N}(n) & \longrightarrow & p^* p_* \mathcal{N}(n-1) & \longrightarrow & p^* p_*(\mathcal{N}(n-1)|_{Y'}) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{N}(n) & \longrightarrow & \mathcal{N}(n-1) & \longrightarrow & \mathcal{N}(n-1)|_{Y'} \longrightarrow 0, \end{array}$$

in which the exactness of the upper row follows from the vanishing of $R^1 p_*(\mathcal{N}(n))$ proved in (i), and note that $p^* p_*(\mathcal{N}(n-1)|_{Y'}) \rightarrow \mathcal{N}(n-1)|_{Y'}$ is surjective for $n \geq 1$ in view of Nakayama's lemma, since its restriction to the fibers of p is surjective by cohomology and base change [44, III, Theorem 12.11].

To prove (iii), we work Zariski-locally around a point $y \in X$. In view of (ii) for $n = 0$, we can assume after shrinking X the existence of a section $\sigma \in H^0(X', \mathcal{N})$ that does not vanish identically on X'_y . Since $\mathcal{N}|_{X'_y} \simeq \mathcal{O}_{X'_y}$ and X'_y is a projective space, the section σ vanishes nowhere on X'_y and, after shrinking X again, it induces an isomorphism $\sigma : \mathcal{O}_{X'} \xrightarrow{\sim} \mathcal{N}$. Assertion (iii) now follows from the fact that the natural morphism $\mathcal{O}_X \rightarrow p_* \mathcal{O}_{X'}$ is an isomorphism [38, VII, Lemme 3.5]. \square

COROLLARY 1.5. – *Under the hypotheses of §1.4.1, if X is moreover a proper variety over k , the formula (1.2) induces an isomorphism of functors*

$$(1.3) \quad \mathrm{Pic}_{X/k} \times \mathbf{Z}_{Y/k} \xrightarrow{\sim} \mathrm{Pic}_{X'/k}.$$

Proof. – Since the formation of the blow-up of a regular closed immersion commutes with flat base change (see [38, VII, Propositions 1.5 et 1.8 i]), we can apply Proposition 1.3 to the morphisms $i_T : Y_T \rightarrow X_T$ for $T \in (\mathrm{Sch}/k)$. \square

1.4.2. *Picard schemes of curves.* – If C is a proper curve over k , then $\mathbf{Pic}_{C/k}$ is smooth over k by [12, 8.4/2]. Moreover, letting $D := \widetilde{C_{k_p}^{\text{red}}}$ be the normalization of the reduction of C_{k_p} , which is a smooth proper curve over k_p , the pull-back morphism $\mathbf{Pic}_{C_{k_p}/k_p}^0 \rightarrow \mathbf{Pic}_{D/k_p}^0$ induces an isomorphism

$$(1.4) \quad A(\mathbf{Pic}_{C_{k_p}/k_p}^0) \xrightarrow{\sim} \mathbf{Pic}_{D/k_p}^0,$$

as [12, 9.2/11] shows. The principal polarization of \mathbf{Pic}_{D/k_p}^0 given by the theta divisor thus induces a canonical principal polarization on $\mathbf{Pic}_{C/k}$ in the sense of §1.2. If C is irreducible, then so is D [35, Proposition 2.4.5] and the principally polarized abelian variety \mathbf{Pic}_{D/k_p}^0 over k_p is thus indecomposable if it is non-zero (see [7, §2.1]).

In preparation for the proof of Proposition 1.6 and, later, for the statement of Corollary 1.8, let us recall that if V is a variety over k and V' denotes the normalization of V^{red} , then V is *geometrically unibranch* if and only if the natural morphism $V' \rightarrow V$ is a universal homeomorphism [35, Proposition 2.4.5, (6.15.1), end of (6.15.3)]. In particular, normal varieties are geometrically unibranch. We also recall that the property of being geometrically unibranch is invariant under extension of scalars [35, Proposition 6.15.7].

PROPOSITION 1.6. – *Let C be a proper curve over k and let $C' := \widetilde{C^{\text{red}}}$ be the normalization of its reduction. Then there is a short exact sequence*

$$(1.5) \quad 0 \rightarrow \mathbf{Pic}_{C/k}^0 \rightarrow \mathbf{Pic}_{C/k} \rightarrow \mathbf{Z}_{C'/k} \rightarrow 0.$$

Proof. – Both $\mathbf{Z}_{C'/k}$ and $\mathbf{Pic}_{C/k} / \mathbf{Pic}_{C/k}^0$ are étale group schemes over k . They are thus isomorphic if and only if so are their base changes $G := \mathbf{Z}_{C'_{k_p}/k_p}$ and $H := \mathbf{Pic}_{C_{k_p}/k_p} / \mathbf{Pic}_{C_{k_p}/k_p}^0$ to k_p . Letting $D := \widetilde{C_{k_p}^{\text{red}}}$ be the normalization of the reduction of C_{k_p} , which is a smooth proper curve over k_p , one has $G = \mathbf{Z}_{D/k_p}$ by Remark 1.2 (ii) (indeed the map $D \rightarrow C'_{k_p}$ is a universal homeomorphism since C'_{k_p} is geometrically unibranch), and $H = \mathbf{Pic}_{D/k_p} / \mathbf{Pic}_{D/k_p}^0$ by [12, 9.2/11]. That $G \simeq H$ now follows from the fact that $G(\bar{k})$ and $H(\bar{k})$ are both isomorphic, as Γ_k -modules, to $\mathbf{Z}(D_{\bar{k}})$. □

1.5. When do Jacobians split?

We now provide, in Theorem 1.7, a criterion for the Jacobian of a proper reduced curve to be the product of an abelian variety and of an affine group scheme. We will use Theorem 1.7 in Lemma 3.8, which plays a key role in the proof of Theorem C.

1.5.1. *Statement.* – Let us introduce some notation. Whenever D is a smooth proper integral curve over k , the *genus* of D is the dimension of the abelian variety $\mathbf{Pic}_{D/k}^0$. We note that D has genus 0 if and only if the irreducible components of $D_{\bar{k}}$ (which are all isomorphic) are rational. Given a proper reduced curve C over k , we denote by C_{rat} (resp. C_{irrat}) the union of those irreducible components B of C such that the normalization of $(B_{k_p})^{\text{red}}$ has genus 0 (resp. has genus ≥ 1). We view C_{rat} and C_{irrat} as reduced closed subschemes of C . We define a *strict cycle of components of $C_{\bar{k}}$* to be a sequence of pairwise distinct irreducible components B_1, \dots, B_n of $C_{\bar{k}}$ for some integer $n \geq 2$, such that there exist pairwise distinct points x_1, \dots, x_n of $C_{\bar{k}}$ with $x_i \in B_i \cap B_{i+1}$ for all $i \in \{1, \dots, n-1\}$ and $x_n \in B_n \cap B_1$.

Finally, we recall that a reduced curve over \bar{k} is *seminormal* if it is étale locally isomorphic to the union of the coordinate axes in an affine space over \bar{k} [59, Chapter I, 7.2.2].

THEOREM 1.7. – *Let C be a proper reduced curve over k . The group scheme $\mathbf{Pic}_{C/k}^0$ is the product of an abelian variety and of an affine group scheme over k if and only if the following conditions all hold:*

- (i) *the scheme $(C_{\text{irrat}})_{\bar{k}}$ is reduced and seminormal and its irreducible components are smooth;*
- (ii) *any strict cycle of components of $C_{\bar{k}}$ is contained in $(C_{\text{rat}})_{\bar{k}}$;*
- (iii) *for every connected component B of C_{rat} , either the scheme $B \cap C_{\text{irrat}}$ is étale over k , or it is of the form $\text{Spec}(k')$ for some field k' and the restriction map $H^0(B, \mathcal{O}_B) \rightarrow k'$ is bijective.*

In this case, the natural map $\mathbf{Pic}_{C/k} \rightarrow \mathbf{Pic}_{C_{\text{rat}}/k} \times \mathbf{Pic}_{C_{\text{irrat}}/k}$ is an isomorphism and $\mathbf{Pic}_{C_{\text{rat}}/k}^0$ is affine while $\mathbf{Pic}_{C_{\text{irrat}}/k}^0$ is an abelian variety.

Condition (iii) holds if $(C_{\text{rat}} \cap C_{\text{irrat}})_{\bar{k}}$ is reduced, and it implies, in turn, that $C_{\text{rat}} \cap C_{\text{irrat}}$ is reduced. The reverse implications are true if k is perfect.

When C is integral and geometrically locally irreducible, for instance when C is integral and geometrically unibranch, Theorem 1.7 takes on a particularly simple form, which we now state. We recall that normal varieties are geometrically unibranch (see §1.4.2 for more reminders on this property). In the sequel, we shall only apply Theorem 1.7 to normal curves, through Corollary 1.8.

COROLLARY 1.8. – *Let C be a proper integral curve over k . Assume that the connected components of $C_{\bar{k}}$ are irreducible; such is the case, for instance, if C is geometrically unibranch. Then the group scheme $\mathbf{Pic}_{C/k}^0$ is the product of an abelian variety and of an affine group scheme over k if and only if at least one of the following two conditions holds:*

- (i) *C is smooth over k (in which case $\mathbf{Pic}_{C/k}^0$ is an abelian variety);*
- (ii) *the normalization D of $(C_{k_p})^{\text{red}}$ has genus 0 (in which case $\mathbf{Pic}_{C/k}^0$ is affine).*

Proof. – Our assumptions imply that $C_{\text{rat}} = \emptyset$ or $C_{\text{irrat}} = \emptyset$, that there is no strict cycle of components of $C_{\bar{k}}$ and that C is smooth over k if and only if $C_{\bar{k}}$ is reduced with smooth irreducible components. Thus, the conditions of Theorem 1.7 are all met if and only if at least one of (i) and (ii) holds. \square

REMARK 1.9. – Corollary 1.8 applies to integral curves that may not be geometrically reduced. It would however fail in general for irreducible but non-reduced curves, as the following example shows. Let E be an elliptic curve over \bar{k} , and let \mathcal{L} be an ample line bundle on E . Define $C := \text{Proj}_E(\mathcal{O}_E \oplus \mathcal{L})$, where sections of \mathcal{L} square to 0. The natural closed immersion $i : E = C^{\text{red}} \rightarrow C$ then induces an isomorphism $i^* : \mathbf{Pic}_{C/k}^0 \xrightarrow{\sim} \mathbf{Pic}_{E/k}^0 \simeq E$. Indeed, in view of (1.4), it suffices to show that the kernel of $(i^*)_{k_p}$ has trivial tangent space at the identity, which follows from the fact that the pull-back $i^* : H^1(C, \mathcal{O}_C) \xrightarrow{\sim} H^1(E, \mathcal{O}_E)$ is an isomorphism. We note that C is geometrically unibranch since C^{red} is normal.

1.5.2. *A few general lemmas.* – We establish a series of lemmas on which the proof of Theorem 1.7 will rely. The first one is due to Tanaka, see [78, Lemma 3.3].

LEMMA 1.10. – *Let F be a field extension of k . If F/k is not separable, there exist finite purely inseparable field extensions $k \subseteq k' \subseteq k''$ and a k' -linear embedding $k'' \hookrightarrow F \otimes_k k'$ such that $F \otimes_k k'$ is a field and $k'' \neq k'$.*

Proof. – Let k'' be a minimal finite purely inseparable field extension of k such that $F \otimes_k k''$ fails to be reduced. Let p denote the characteristic of k . As k''/k is finite, purely inseparable and non-trivial, there exists a subextension k'/k such that $[k'' : k'] = p$. Fix $x \in k'$ such that $k'' = k'(x^{1/p})$. By the minimality of k'' , the finite connected non-zero F -algebra $F' = F \otimes_k k'$ is reduced, hence is a field. On the other hand, as $F \otimes_k k'' = F'[t]/(t^p - x)$ is not reduced, we see that x must be a p -th power in F' , so that k'' embeds k' -linearly into F' . \square

LEMMA 1.11. – *Let $f : G \rightarrow G'$ be a surjective morphism between connected smooth group schemes over k such that the kernel of $A(f_{k_p}) : A(G_{k_p}) \rightarrow A(G'_{k_p})$ is smooth and connected. If G is isomorphic to the product of an abelian variety and of an affine group scheme over k , then so is G' .*

Proof. – Suppose $G = L \times A$ with L affine and A an abelian variety. Let $K = \text{Ker}(f)$. Let $p : G \rightarrow A$ be the projection, let $i : A \rightarrow G$ be the inclusion, and let $p(K)$ denote the scheme-theoretic image of K by p ; it is a subgroup scheme of A [43, VI_A, Proposition 6.4]. We note that $A(G_{k_p}) = A_{k_p}$ and that $p(K)_{k_p} \subseteq \text{Ker}(A(f_{k_p}))$. The morphism $\text{Ker}(A(f_{k_p})) \rightarrow G'_{k_p}$ induced by $f \circ i$ factors through $L(G'_{k_p})$ and hence vanishes since $\text{Ker}(A(f_{k_p}))$ is an abelian variety and $L(G'_{k_p})$ is affine. Therefore the morphism $p(K) \rightarrow G'$ induced by $f \circ i$ also vanishes. We deduce that $i(p(K)) \subseteq K$, hence $K = (K \cap L) \times i(p(K))$. On the other hand, the closed immersion $G/K \rightarrow G'$ induced by f is an isomorphism as it is surjective and G' is reduced. It follows that $G' = (L/(K \cap L)) \times (A/p(K))$, and the lemma is proved. \square

LEMMA 1.12. – *Let C be a proper reduced curve over k and B be a geometrically connected reduced closed subscheme of C of pure dimension 1. View the union of the irreducible components of C not contained in B as a reduced closed subscheme B' of C . If the natural morphism $B \cap B' \rightarrow \text{Spec}(H^0(B', \mathcal{O}_{B'}))$ is étale and if any strict cycle of components of $C_{\bar{k}}$ is contained in $B'_{\bar{k}}$, then the natural morphism $\mathbf{Pic}_{C/k} \rightarrow \mathbf{Pic}_{B/k} \times \mathbf{Pic}_{B'/k}$ is an isomorphism.*

Proof. – If $i : B_T \hookrightarrow C_T, i' : B'_T \hookrightarrow C_T, i'' : B_T \cap B'_T \hookrightarrow C_T$ denote the inclusions, the short exact sequence $1 \rightarrow \mathbf{G}_m \rightarrow i_* \mathbf{G}_m \times i'_* \mathbf{G}_m \rightarrow i''_* \mathbf{G}_m \rightarrow 1$ of sheaves for the Zariski topology on C_T induces, for any $T \in (\text{Sch}/k)$, an exact sequence of groups

$$(1.6) \quad \begin{aligned} 1 \rightarrow \mathbf{G}_m(C_T) &\rightarrow \mathbf{G}_m(B_T) \times \mathbf{G}_m(B'_T) \rightarrow \mathbf{G}_m(B_T \cap B'_T) \\ &\rightarrow \text{Pic}(C_T) \rightarrow \text{Pic}(B_T) \times \text{Pic}(B'_T) \rightarrow \text{Pic}(B_T \cap B'_T). \end{aligned}$$

The natural morphism $B \cap B' \rightarrow \text{Spec}(H^0(B', \mathcal{O}_{B'}))$ is an étale morphism between finite k -schemes that induces an injection on \bar{k} -points, in view of the assumption about strict cycles of components. (Note that $\text{Spec}(H^0(B', \mathcal{O}_{B'}))(\bar{k})$ is the set of connected components of $B'_{\bar{k}}$.)

It is therefore an open and closed immersion [41, I, Théorème 5.1]. In particular, the resulting restriction map

$$\mathbf{G}_m(\mathrm{Spec}(H^0(B', \mathcal{O}_{B'})) \times_k T) \rightarrow \mathbf{G}_m(B_T \cap B'_T)$$

is onto, and hence so is the restriction map $\mathbf{G}_m(B'_T) \rightarrow \mathbf{G}_m(B_T \cap B'_T)$. Noting that $\mathbf{Pic}_{B \cap B'/k} = 0$, the conclusion of the lemma now results from the exact sequence of fppf sheaves obtained by sheafifying (1.6) with respect to T . \square

LEMMA 1.13. – *Let k' be a finite purely inseparable extension of k . Let C be a proper integral curve over k' . We view C as a curve over k . If $\mathbf{Pic}_{C/k}^0$ is the product of an abelian variety and of an affine group scheme over k , then $k' = k$ or $C = C_{\mathrm{rat}}$.*

Proof. – For all $T \in (\mathrm{Sch}/k)$, pull-back induces an equivalence between the categories of étale T -schemes and of étale $T_{k'}$ -schemes [41, IX, Théorème 4.10]. It follows at once that the natural morphism

$$\mathrm{Res}_{k'/k}(\mathbf{Pic}_{C/k'}) \rightarrow \mathrm{Res}_{k'/k}(\mathbf{Pic}_{C/k', \acute{e}t})$$

of functors $(\mathrm{Sch}/k)^{\mathrm{op}} \rightarrow (\mathrm{Ab})$ becomes an isomorphism after étale sheafification. On the other hand, by the very definition of the absolute Picard functor (see §1.4) and of Weil restriction of scalars, one has $\mathbf{Pic}_{C/k} = \mathrm{Res}_{k'/k}(\mathbf{Pic}_{C/k'})$. By these two remarks, we get, after sheafification, an isomorphism $\mathbf{Pic}_{C/k} = \mathrm{Res}_{k'/k}(\mathbf{Pic}_{C/k'})$, which restricts to an isomorphism $\mathbf{Pic}_{C/k}^0 = \mathrm{Res}_{k'/k}(\mathbf{Pic}_{C/k'}^0)$ in view of [19, Proposition A.5.9]. Let us write $\mathbf{Pic}_{C/k}^0 = L \times A$ with L affine and A an abelian variety. Applying [37, XVII, Appendice III, Proposition 5.1] with $U = L$ (or [19, Proposition A.7.8]) now shows that if $k' \neq k$, then $\mathbf{Pic}_{C/k}^0 = L$, so that $A((\mathbf{Pic}_{C/k}^0)_{k_p}) = 0$ and $C = C_{\mathrm{rat}}$. \square

LEMMA 1.14. – *Let D be the disjoint union of smooth proper geometrically integral curves D_1, \dots, D_n over k . Let C be a proper curve over k such that $H^0(C, \mathcal{O}_C) = k$ and $C(k) \neq \emptyset$. Let $v : D \rightarrow C$ be a morphism. If $v^* : \mathbf{Pic}_{C/k} \rightarrow \mathbf{Pic}_{D/k}$ admits a section, then there exist morphisms $\rho_i : C \rightarrow \mathbf{Pic}_{D_i/k}$ for $i \in \{1, \dots, n\}$ such that $\rho_i \circ v|_{D_i}$ is the canonical morphism for all i while $\rho_i \circ v|_{D_j}$ is constant for all $j \neq i$.*

Proof. – We let $v_i : D_i \rightarrow C$ be the restriction of v , fix a section $\sigma : \mathbf{Pic}_{D/k} \rightarrow \mathbf{Pic}_{C/k}$ of v^* and, noting that $\mathbf{Pic}_{D/k} = \prod_{i=1}^n \mathbf{Pic}_{D_i/k}$, let $\sigma_i : \mathbf{Pic}_{D_i/k} \rightarrow \mathbf{Pic}_{C/k}$ be the restriction of σ , so that $v_i^* \circ \sigma_j = \delta_{ij}$. Let $\iota_i : D_i \rightarrow \mathbf{Pic}_{D_i/k}$ be the canonical morphism.

Then v_j^* maps $\sigma_i \circ \iota_i \in \mathbf{Pic}_{C/k}(D_i)$ to $\iota_i \in \mathbf{Pic}_{D_i/k}(D_i)$ if $j = i$, to $0 \in \mathbf{Pic}_{D_j/k}(D_i)$ otherwise. As $H^0(C, \mathcal{O}_C) = H^0(D_i, \mathcal{O}_{D_i}) = k$ and $C(k) \neq \emptyset$, there are a canonical bijection $\mathrm{Pic}(C_T)/\mathrm{Pic}(T) \xrightarrow{\sim} \mathbf{Pic}_{C/k}(T)$ and a canonical injection $\mathrm{Pic}((D_i)_T)/\mathrm{Pic}(T) \hookrightarrow \mathbf{Pic}_{D_i/k}(T)$ for all T . We deduce, for each i , the existence of $\alpha_i \in \mathrm{Pic}(C \times_k D_i)$ such that $(v_j \times 1)^* \alpha_i \in \mathrm{Pic}(D_j \times_k D_i)$ is the class of the diagonal for $j = i$ and comes by pull-back from $\mathrm{Pic}(D_i)$ for all $j \neq i$. Switching the factors, the class α_i gives rise to the desired element $\rho_i \in \mathbf{Pic}_{D_i/k}(C)$. \square

LEMMA 1.15. – *Let C be a proper curve over k . Let $v : C' \rightarrow C$ be the normalization of C^{red} . The pull-back morphism $v^* : \mathbf{Pic}_{C/k}^0 \rightarrow \mathbf{Pic}_{C'/k}^0$ is surjective.*

Proof. – We may assume that the field k is separably closed. We then claim that the map $v^* : \mathbf{Pic}_{C/k}(k) \rightarrow \mathbf{Pic}_{C'/k}(k)$ is onto. As $\mathbf{Pic}_{C'/k}$ is smooth over k , it will follow that the morphism $v^* : \mathbf{Pic}_{C/k} \rightarrow \mathbf{Pic}_{C'/k}$ is dominant, by [12, 2.2/13], so that the morphism $v^* : \mathbf{Pic}_{C/k}^0 \rightarrow \mathbf{Pic}_{C'/k}^0$ is dominant, by Proposition 1.6, and hence surjective, by [43, VI_A, Corollaire 6.2 (i)]. To prove the claim, we may assume that C is reduced, by [12, 9.2/5]. Letting $C^0 \subseteq C$ be a dense open normal subset, we then remark that $v^* : \mathrm{Pic}(C) \rightarrow \mathrm{Pic}(C')$ is onto since any divisor on C' is linearly equivalent to a divisor supported on $v^{-1}(C^0)$. As k is separably closed, it follows that $v^* : \mathrm{Pic}_{C/k,\acute{e}t}(k) \rightarrow \mathrm{Pic}_{C'/k,\acute{e}t}(k)$ is onto as well, as desired. \square

1.5.3. *Proof of Theorem 1.7.* – As the formation of C_{rat} and C_{irrat} is compatible with separable extensions of scalars and as the assertions of the last sentence of the theorem are of a geometric nature, we may, and will henceforth, assume that k is separably closed. For later use, we note that thanks to this assumption, if condition (i) of Theorem 1.7 holds, then the irreducible components of C_{irrat} , viewed as reduced schemes, are geometrically reduced (being both reduced and generically geometrically reduced) and therefore smooth over k , and the non-smooth locus of C_{irrat} over k consists of k -points (as the intersection of any two irreducible components of C_{irrat} is étale).

Step 1. – We assume that (i)–(iii) hold and deduce the remaining assertions.

From (1.4) applied to C_{rat} , we deduce that the group scheme $(\mathbf{Pic}_{C_{\mathrm{rat}}/k}^0)_{k_p}$ is affine; hence $\mathbf{Pic}_{C_{\mathrm{rat}}/k}^0$ is also affine. To prove that $\mathbf{Pic}_{C_{\mathrm{irrat}}/k}^0$ is an abelian variety and that the natural map $\mathbf{Pic}_{C/k} \rightarrow \mathbf{Pic}_{C_{\mathrm{rat}}/k} \times \mathbf{Pic}_{C_{\mathrm{irrat}}/k}$ is an isomorphism, we argue by induction on the number of irreducible components of C_{irrat} . When $C_{\mathrm{irrat}} = \emptyset$, there is nothing to prove. Otherwise, let us choose an irreducible component B of C_{irrat} and denote by B' (resp. B'_{irrat}) the union of the irreducible components of C (resp. of C_{irrat}) that are distinct from B . We view B , B' and B'_{irrat} as reduced closed subschemes of C . The equalities $C = B \cup B' = C_{\mathrm{rat}} \cup C_{\mathrm{irrat}} = B \cup C_{\mathrm{rat}} \cup B'_{\mathrm{irrat}}$ and $B' = C_{\mathrm{rat}} \cup B'_{\mathrm{irrat}}$ induce a commutative square

$$(1.7) \quad \begin{array}{ccc} \mathbf{Pic}_{C/k} & \longrightarrow & \mathbf{Pic}_{B/k} \times \mathbf{Pic}_{B'/k} \\ \downarrow & & \downarrow \wr \\ \mathbf{Pic}_{C_{\mathrm{rat}}/k} \times \mathbf{Pic}_{C_{\mathrm{irrat}}/k} & \longrightarrow & \mathbf{Pic}_{B/k} \times \mathbf{Pic}_{C_{\mathrm{rat}}/k} \times \mathbf{Pic}_{B'_{\mathrm{irrat}}/k} \end{array}$$

whose right vertical arrow is an isomorphism by the induction hypothesis. To conclude the proof, we need only verify that Lemma 1.12 can be applied, on the one hand, to C , B and B' , and on the other hand, to C_{irrat} , B and B'_{irrat} . Indeed, it will follow that the horizontal arrows of (1.7) are isomorphisms, which implies, on the one hand, that the left vertical arrow of this square is an isomorphism, and on the other hand, that $\mathbf{Pic}_{C_{\mathrm{irrat}}/k}^0$ is an abelian variety, since so are $\mathbf{Pic}_{B'_{\mathrm{irrat}}/k}^0$ (by the induction hypothesis) and $\mathbf{Pic}_{B/k}^0$ (as B is smooth over k [12, 9.2/3]).

To this end, letting F stand either for B' or for B'_{irrat} , we fix $x \in B \cap F$, and denote by E the connected component of x in F and by C_1, \dots, C_m the irreducible components of C_{irrat} containing x , numbered so that $B = C_1$. The finite k -algebra $H^0(E, \mathcal{O}_E)$, being non-zero, connected and reduced, is a field; it embeds into the residue field $k(x)$ of x . We now have to prove that the natural morphism $B \cap E \rightarrow \mathrm{Spec}(H^0(E, \mathcal{O}_E))$ is étale at x .

If $k(x) \neq k$, then $m = 1$ and therefore $C_{\text{rat}} \cap C_{\text{irrat}}$ is not étale over k at x . We deduce from (iii) that E coincides with the connected component of x in C_{rat} , that $B \cap E$ is reduced at x and that $H^0(E, \mathcal{O}_E) = k(x)$; hence the desired result.

If $k(x) = k$, then $H^0(E, \mathcal{O}_E) = k$ and it suffices to see that $B \cap E$ is reduced at x . In the Zariski tangent space $T_x C$ of C at x , we have $T_x E = V + T_x C_2 + \dots + T_x C_m$ where $V = T_x C_{\text{rat}}$ if $x \in C_{\text{rat}}$ and $F = B'$, and $V = 0$ otherwise. It follows from (iii) that $C_{\text{rat}} \cap C_{\text{irrat}}$ is reduced, so that $V \cap (T_x C_1 + \dots + T_x C_m) = 0$, and from (i) that the lines $T_x C_1, \dots, T_x C_m$ are in direct sum. Hence $T_x(B \cap E) = T_x C_1 \cap T_x E = 0$, and $B \cap E$ is indeed reduced at x . Step 1 is complete.

We now assume that $\mathbf{Pic}_{C/k}^0$ is the product of an abelian variety and of an affine group scheme over k , and prove (i)–(iii) in four more steps. By Lemma 1.11, we may assume that C is connected. In addition, we may assume that $C_{\text{irrat}} \neq \emptyset$. Let C_1, \dots, C_n be the irreducible components of C_{irrat} , viewed as reduced schemes. Let D_i be the normalization of C_i , let D be the disjoint union of the D_i and let $v_i : D_i \rightarrow C$ and $v : D \rightarrow C$ be the natural morphisms.

Step 2. – We prove that C_1, \dots, C_n and C_{irrat} are geometrically reduced over k .

As $C_{\text{irrat}} = C_1 \cup \dots \cup C_n$, it suffices to check that the C_i are geometrically reduced. Assume that some $B \in \{C_1, \dots, C_n\}$ is not geometrically reduced. By Lemma 1.10 applied to $k(B)$, there exist subfields $k \subseteq k' \subseteq k'' \subseteq k_p$, with $k'' \neq k'$ and k''/k finite, such that $B_{k'}$ is integral and such that if B' denotes its normalization, the natural morphism $B' \rightarrow \text{Spec}(k')$ factors through $\text{Spec}(k'')$. Let $\omega : C' \rightarrow C_{k'}$ be the normalization of $(C_{k'})^{\text{red}}$. By Lemma 1.15, the pull-back map $\omega^* : \mathbf{Pic}_{C_{k'}/k'}^0 \rightarrow \mathbf{Pic}_{C'/k'}^0$ is surjective, and (1.4) shows that $A(\omega_{k_p}^*)$ is an isomorphism. As B' is a connected component of C' , we deduce, thanks to Lemma 1.11, that $\mathbf{Pic}_{B'/k'}^0$ is the product of an abelian variety and of an affine group scheme over k' . Lemma 1.13 implies that $k'' = k'$ or $B' = B'_{\text{rat}}$, a contradiction.

Step 3. – Assuming that D is smooth over k , we construct $\rho_i : C \rightarrow \mathbf{Pic}_{D_i/k}^1$ such that $\rho_i \circ v_i$ is a closed immersion while $\rho_i(C_{\text{rat}} \cup \bigcup_{j \neq i} C_j)$ is finite, for all i .

It suffices to check that the hypotheses of Lemma 1.14 are satisfied. Indeed, the canonical morphism $D_i \rightarrow \mathbf{Pic}_{D_i/k}^1$ is a closed immersion if D_i is a smooth proper integral curve of genus ≥ 1 over k [69, Propositions 6.1 and 2.3], and $\rho_i(C_{\text{rat}})$ is finite since any morphism from a rational curve to an abelian variety is constant.

We recall that $C_{\text{irrat}} \neq \emptyset$. As C is proper, connected and reduced, the restriction map $H^0(C, \mathcal{O}_C) \rightarrow H^0(C_1, \mathcal{O}_{C_1})$ has to be injective; as C_1 is geometrically integral, we deduce that $H^0(C, \mathcal{O}_C) = k$. In addition, as k is separably closed, we have $C_1(k) \neq \emptyset$ [12, 2.2/13], hence $C(k) \neq \emptyset$.

As D is smooth, the group scheme $\mathbf{Pic}_{D/k}^0$ is an abelian variety and the morphism $v^* : \mathbf{Pic}_{C/k}^0 \rightarrow \mathbf{Pic}_{D/k}^0$ induces an isomorphism $A((\mathbf{Pic}_{C/k}^0)_{k_p}) \xrightarrow{\sim} (\mathbf{Pic}_{D/k}^0)_{k_p}$ (see (1.4)). Our assumption on $\mathbf{Pic}_{C/k}^0$ therefore implies that $v^* : \mathbf{Pic}_{C/k}^0 \rightarrow \mathbf{Pic}_{D/k}^0$ admits a section. Letting C' be the normalization of C^{red} , the natural map $v^* : \mathbf{Z}_{C'/k} \rightarrow \mathbf{Z}_{D/k}$ also admits a section. In addition, the sequence (1.5) splits as k is separably closed and $\mathbf{Pic}_{C/k}^0$ is smooth, and the choice of a splitting of (1.5) and of a section of $v^* : \mathbf{Z}_{C'/k} \rightarrow \mathbf{Z}_{D/k}$ determines a splitting of the sequence (1.5) associated with D . Applying Proposition 1.6 to C and to D , we now conclude that $v^* : \mathbf{Pic}_{C/k} \rightarrow \mathbf{Pic}_{D/k}$ admits a section. This completes Step 3.

Step 4. – We prove (i) and (ii) of Theorem 1.7.

Step 2 ensures that $((C_{\bar{k}})^{\text{red}})_{\text{irrat}} = (C_{\text{irrat}})_{\bar{k}}$ and it follows from Lemma 1.11, in view of [12, 9.2/5], that $\mathbf{Pic}_{(C_{\bar{k}})^{\text{red}}/\bar{k}}^0$ is the product of an abelian variety and of an affine group scheme over \bar{k} . Thus, after replacing C with $(C_{\bar{k}})^{\text{red}}$, we may, and will until the end of Step 4, assume that $k = \bar{k}$. In this case D is smooth over k and Step 3 becomes applicable.

To complete the proof of (i), it suffices to show that for all $i \in \{1, \dots, n\}$, the curve C_i is smooth and the scheme $C_i \cap (\bigcup_{j \neq i} C_j)$ is reduced. We fix i . As $\rho_i \circ v_i$ is an immersion, so is v_i ; as $v_i(D_i) = C_i$ is a reduced closed subscheme of C , we deduce that v_i restricts to an isomorphism $D_i \rightarrow C_i$, so that C_i is smooth. Now the restriction of ρ_i to the subscheme $C_i \cap (\bigcup_{j \neq i} C_j)$ is both a closed immersion (since so is $\rho_i|_{C_i}$) and a morphism whose scheme-theoretic image is finite and reduced (since so is $\rho_i|_{\bigcup_{j \neq i} C_j}$), hence this scheme is reduced.

To prove (ii), we pick a strict cycle of components B_1, \dots, B_n of C and pairwise distinct points $x_i \in B_i \cap B_{i+1}$ for $i \in \{1, \dots, n-1\}$ and $x_n \in B_n \cap B_1$. If one of the B_i were contained in C_{irrat} , say $B_1 = C_1$, then $\rho_1(x_1)$ and $\rho_1(x_n)$ would have to be distinct, since $\rho_1|_{C_1}$ is injective, and equal, since $\rho_1(B_2 \cup \dots \cup B_n)$ is a point (being finite and connected). This is absurd.

Step 5. – We prove (iii) of Theorem 1.7.

We now know that (i) holds, hence D is smooth over k : we can apply Step 3 again.

LEMMA 1.16. – *Let $i \in \{1, \dots, n\}$. For any purely 1-dimensional connected reduced closed subscheme E of C such that $E \cap C_i$ is finite and non-empty, the restriction map $H^0(E, \mathcal{O}_E) \rightarrow H^0(E \cap C_i, \mathcal{O}_{E \cap C_i})$ is an isomorphism of fields.*

Proof. – The morphism $\rho_i|_E$ has finite image (by Step 3), hence it factors through an affine open $\text{Spec}(R) \subset \mathbf{Pic}_{D_i/k}^1$. To see that the map of the lemma is surjective, we note that its composition with $R \rightarrow H^0(E, \mathcal{O}_E)$ is surjective as $\rho_i|_{E \cap C_i}$ is a closed immersion. It is also injective, as $H^0(E, \mathcal{O}_E)$ is a field (being a non-zero, connected, reduced, finite k -algebra) and $E \cap C_i \neq \emptyset$. \square

To prove (iii), let B be a connected component of C_{rat} such that $B \cap C_{\text{irrat}}$ is not étale over k , say at a point x .

After renumbering, we may assume that the C_i containing x are C_1, \dots, C_m , for some $m \in \{1, \dots, n\}$. If x were a k -point, the subspace $T_x B \cap T_x C_{\text{irrat}}$ of the Zariski tangent space $T_x C$ would be non-zero. After renumbering C_1, \dots, C_m appropriately and setting $E = B \cup C_2 \cup \dots \cup C_m$, the vector space $T_x E \cap T_x C_1$ would be non-zero. The scheme $E \cap C_1$ would then be non-reduced; this would contradict Lemma 1.16. Thus $k(x) \neq k$ and hence $m = 1$.

Let E' be the connected component of x in $C_{\text{rat}} \cup C_2 \cup \dots \cup C_n$.

As the evaluation map $H^0(E' \cap C_1, \mathcal{O}_{E' \cap C_1}) \rightarrow k(x)$ is surjective, Lemma 1.16 shows that $H^0(E', \mathcal{O}_{E'}) = H^0(E' \cap C_1, \mathcal{O}_{E' \cap C_1}) = k(x)$; hence, by Step 2, there is no k -algebra morphism $H^0(E', \mathcal{O}_{E'}) \rightarrow H^0(C_j, \mathcal{O}_{C_j})$ for any j . Therefore $E' = B$ and $B \cap C_{\text{irrat}} = E' \cap C_1$. This completes Step 5 as well as the proof of Theorem 1.7.

1.6. Murre's intermediate Jacobian

If X is a smooth projective variety over k , Murre defines an *algebraic representative for algebraically trivial codimension 2 cycles on $X_{\bar{k}}$* to be an abelian variety $\text{Ab}^2(X_{\bar{k}})$ over \bar{k} endowed with a morphism

$$(1.8) \quad \phi_X^2 : \text{CH}^2(X_{\bar{k}})_{\text{alg}} \rightarrow \text{Ab}^2(X_{\bar{k}})(\bar{k})$$

that is initial among regular homomorphisms with values in an abelian variety (see [72, Definition 1.6.1, §1.8]). It is obviously unique up to a unique isomorphism, Murre has shown its existence (see [72, Theorem A p. 226] and [54]), and Achter, Casalaina-Martin and Vial have shown that it descends uniquely to an abelian variety $\text{Ab}^2(X_{k_p})$ over k_p in such a way that (1.8) is Γ_k -equivariant [3, Theorem 4.4]. If X is a smooth projective \bar{k} -rational threefold over k , we endow $\text{Ab}^2(X_{k_p})$ with the principal polarization $\theta_X \in \text{NS}^1(\text{Ab}^2(X_{\bar{k}}))^{\Gamma_k}$ constructed in [7, Property 2.4, Corollary 2.8].

1.7. Representability lemmas

Here are three lemmas to be used later.

LEMMA 1.17. – *Let F and F' be two functors $(\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$. If $F \times F'$ is represented by a group scheme locally of finite type over k , then so is F .*

Proof. – Let G be the group scheme represented by $F \times F'$ and let $\mu : G \rightarrow G$ be the morphism induced by the endomorphism $(x, y) \mapsto (0, y)$ of $F \times F'$. Then F is represented by $\text{Ker}(\mu)$, which is a group scheme locally of finite type over k . \square

LEMMA 1.18. – *Let G be a commutative group scheme locally of finite type over k . Let $v : \mathbf{Z} \rightarrow G$ be a morphism of k -group schemes such that $v(n) \notin G^0(k)$ if $n \neq 0$. Then the cokernel functor $Q : (\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$ defined by $Q(T) = G(T)/v(\mathbf{Z}(T))$ is represented by a group scheme locally of finite type over k .*

Proof. – Translation by $v(n)$ for $n \in \mathbf{Z}$ induces an action of the group \mathbf{Z} on the set of connected components of G . Choose one connected component of G in each orbit of this action. Their disjoint union represents Q . \square

LEMMA 1.19. – *Let $F : (\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$ be a functor and k' be a finite extension of k . Let $\tau \in \{\text{ét}, \text{fppf}\}$. Assume that k'/k is separable if $\tau = \text{ét}$. If F is a sheaf for the topology τ and the functor $(\text{Sch}/k')^{\text{op}} \rightarrow (\text{Ab})$ obtained by restricting F is represented by a group scheme locally of finite type over k' , then F is represented by a group scheme locally of finite type over k .*

Proof. – Let $F' : (\text{Sch}/k')^{\text{op}} \rightarrow (\text{Ab})$ denote the restriction of F and G' the k -group scheme that represents F' . As F' is the restriction of F , there is a canonical descent datum on F' with respect to $p : \text{Spec}(k') \rightarrow \text{Spec}(k)$. This descent datum induces a descent datum on G' . By [70, Lemma I.8.6], the latter is effective, and G' descends to a group scheme G locally of finite type over k . It remains to note that F and $\Phi_G : (\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$ are two τ -sheaves, that p is a τ -covering and that there is by construction an isomorphism of τ -sheaves $p^*F \simeq p^*\Phi_G$ that satisfies the cocycle condition; from this, it follows that $F \simeq \Phi_G$. \square

2. *K*-theory functors

We now define and study several functors $(\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$ built from *K*-theory. Our goal, met in §2.3.2, is to define, for a smooth projective \bar{k} -rational threefold *X* over *k*, a functor $\text{CH}_{X/k, \text{fppf}}^2 : (\text{Sch}/k)^{\text{op}} \rightarrow (\text{Ab})$ that will serve as a substitute for its intermediate Jacobian.

2.1. *K*-theory

We follow the conventions of [38, 80].

2.1.1. *Definition.* – If *X* is a qcqs scheme, we let $\mathbf{K}_0(X)$ be the Grothendieck group of the triangulated category of perfect complexes of \mathcal{O}_X -modules. (This group is denoted by $\mathbf{K}^*(X)$ in [38, IV, Définition 2.2] and coincides with the one defined in [80, §3.1], as indicated in [80, §3.1.1].) We endow $\mathbf{K}_0(X)$ with the ring structure induced by the tensor product ([38, IV, §2.7 b]), [80, §3.15]).

If *X* admits an ample family of line bundles [80, §2.1.1] (for instance, if *X* is quasi-projective over an affine scheme [80, §2.1.2 (c)]), then $\mathbf{K}_0(X)$ is naturally isomorphic to the Grothendieck group of the exact category of vector bundles on *X* (combine [80, Corollary 3.9] and [80, Theorem 1.11.7]).

2.1.2. *Functoriality.* – Let $f : X \rightarrow Y$ be a morphism of qcqs schemes.

The derived pull-back Lf^* of perfect complexes along *f* induces a morphism $f^* : \mathbf{K}_0(Y) \rightarrow \mathbf{K}_0(X)$ ([38, IV, §2.7 b]), [80, §3.14]).

The derived push-forward Rf_* induces a morphism $f_* : \mathbf{K}_0(X) \rightarrow \mathbf{K}_0(Y)$ if Rf_* preserves perfect complexes [80, §3.16]. This condition is satisfied if *f* is a proper and perfect morphism [65, Proposition 2.1, Example 2.2 (a)], for instance if *f* is a proper lci morphism (see [38, VIII, Proposition 1.7]). In this case, the projection formula [38, IV, (2.12.4)] stemming from [38, III, Proposition 3.7] shows that

$$(2.1) \quad f_*(x \otimes f^*y) = f_*x \otimes y$$

for all $x \in \mathbf{K}_0(X)$ and $y \in \mathbf{K}_0(Y)$.

2.1.3. *Rank and determinant.* – The rank $\text{rk} : \mathbf{K}_0(X) \rightarrow \mathbf{Z}(X)$ and the determinant $\det : \mathbf{K}_0(X) \rightarrow \text{Pic}(X)$ are group homomorphisms that are functorial with respect to pull-backs and are such that if $x \in \mathbf{K}_0(X)$ is represented by a bounded complex of vector bundles on *X*, then $\text{rk}(x)$ is the alternating sum of the ranks of its terms and $\det(x)$ is the alternating tensor product of the determinants of its terms (see [58, Theorem 2 p. 42]).

We define $\text{SK}_0(X)$ to be the kernel of $(\text{rk}, \det) : \mathbf{K}_0(X) \rightarrow \mathbf{Z}(X) \times \text{Pic}(X)$.

2.1.4. *Projective bundles and blow-ups.* – If *X* is a qcqs scheme and $\pi : \mathbf{P}_X \mathcal{E} \rightarrow X$ is the projective bundle associated with a vector bundle \mathcal{E} of rank *r* on *X*, the morphism $\mathbf{K}_0(X)^r \rightarrow \mathbf{K}_0(\mathbf{P}_X \mathcal{E})$ given by the formula

$$(2.2) \quad (x_0, \dots, x_{r-1}) \mapsto \sum_{j=0}^{r-1} \pi^* x_j \otimes [\mathcal{O}_{\mathbf{P}_X \mathcal{E}}(-j)]$$

is an isomorphism of $\mathbf{K}_0(X)$ -modules ([38, VI, Théorème 1.1], [80, Theorem 4.1]).

If $i : Y \rightarrow X$ is a regular closed immersion of qcqs schemes of pure codimension $c \geq 1$, if $p : X' \rightarrow X$ is the blow-up of X along Y with exceptional divisor Y' , and if we denote by $p' : Y' \rightarrow Y$ and $i' : Y' \rightarrow X'$ the natural morphisms, Thomason has shown that the morphism $\mathbf{K}_0(X) \times \mathbf{K}_0(Y)^{c-1} \rightarrow \mathbf{K}_0(X')$ given by the formula

$$(2.3) \quad (x, y_1, \dots, y_{c-1}) \mapsto p^*x + \sum_{j=1}^{c-1} i'_* p'^* y_j \otimes [\mathcal{O}_{X'}(jY')]$$

is an isomorphism of $\mathbf{K}_0(X)$ -modules [79, Théorème 2.1].

2.1.5. *Coherent sheaves.* – If X is a qcqs scheme, we let $G_0(X)$ be the Grothendieck group of the triangulated category of pseudo-coherent complexes of \mathcal{O}_X -modules with bounded cohomology. (This group is denoted by $K_*(X)$ in [38, IV, Définition 2.2], see [80, §3.3].)

If X is noetherian, the group $G_0(X)$ is naturally isomorphic to the Grothendieck group of the abelian category of coherent \mathcal{O}_X -modules [38, IV, §2.4]. In this case, letting $F_d G_0(X) \subseteq G_0(X)$ be the subgroup generated by classes of coherent sheaves whose support has dimension $\leq d$ [38, X, Définition 1.1.1] defines a filtration F_\bullet on $G_0(X)$.

If X is noetherian and regular, the natural morphism $\mathbf{K}_0(X) \rightarrow G_0(X)$ is an isomorphism [80, Theorem 3.21]. This allows one to speak of the class of a coherent \mathcal{O}_X -module in $\mathbf{K}_0(X)$, and of the filtration $F_\bullet \mathbf{K}_0(X)$ of $\mathbf{K}_0(X)$ by dimension of the support.

If X is a smooth variety of pure dimension n over k , we use the notation $F^c \mathbf{K}_0(X) = F_{n-c} \mathbf{K}_0(X)$ and $\mathrm{Gr}_F^c \mathbf{K}_0(X) = F^c \mathbf{K}_0(X) / F^{c+1} \mathbf{K}_0(X)$. According to [31, Example 15.1.5], associating with an integral closed subscheme $Z \subseteq X$ of codimension c the class $[\mathcal{O}_Z] \in \mathbf{K}_0(X)$ induces a surjective morphism

$$(2.4) \quad \varphi^c : \mathrm{CH}^c(X) \rightarrow \mathrm{Gr}_F^c \mathbf{K}_0(X).$$

As explained in [31, Example 15.3.6], Jouanolou’s Riemann-Roch theorem without denominators [53] shows that φ^0, φ^1 and φ^2 are isomorphisms, with inverses given by the rank rk , the determinant \det and the opposite $-c_2$ of the second Chern class. In particular, $F^2 \mathbf{K}_0(X) = \mathrm{SK}_0(X)$.

LEMMA 2.1. – *Let $f : X \rightarrow Y$ be a morphism of smooth equidimensional varieties over k . There exists a commutative diagram*

$$(2.5) \quad \begin{array}{ccc} \mathrm{CH}^2(Y) & \xrightarrow[\sim]{\varphi^2} & \mathrm{Gr}_F^2 \mathbf{K}_0(Y) \\ \downarrow f^* & & \downarrow \\ \mathrm{CH}^2(X) & \xrightarrow[\sim]{\varphi^2} & \mathrm{Gr}_F^2 \mathbf{K}_0(X) \end{array}$$

in which the right vertical arrow is induced by $f^* : \mathbf{K}_0(Y) \rightarrow \mathbf{K}_0(X)$.

Proof. – The pull-back $f^* : \mathbf{K}_0(Y) \rightarrow \mathbf{K}_0(X)$ restricts to $f^* : \mathrm{SK}_0(Y) \rightarrow \mathrm{SK}_0(X)$, hence induces $f^* : F^2 \mathbf{K}_0(Y) \rightarrow F^2 \mathbf{K}_0(X)$. Since the morphisms φ^2 are bijective with inverse given by $-c_2$, and since the second Chern class is functorial, we see that $f^* : F^2 \mathbf{K}_0(Y) \rightarrow F^2 \mathbf{K}_0(X)$ induces a morphism $s f^* : \mathrm{Gr}_F^2 \mathbf{K}_0(Y) \rightarrow \mathrm{Gr}_F^2 \mathbf{K}_0(X)$ making the diagram (2.5) commute. \square

2.2. The functor $K_{0,X/k}$ and its sheafifications

2.2.1. *Definition.* – If X is a proper variety over k , the *absolute K_0 functor* of X is

$$\begin{aligned} K_{0,X/k} : (\text{Sch}/k)^{\text{op}} &\rightarrow (\text{Ab}) \\ T &\mapsto K_0(X_T). \end{aligned}$$

The rank and the determinant (see §2.1.3) give rise to morphisms of functors $\text{rk} : K_{0,X/k} \rightarrow \mathbf{Z}_{X/k}$ and $\text{det} : K_{0,X/k} \rightarrow \text{Pic}_{X/k}$. We let $\text{SK}_{0,X/k}$ be the kernel of $(\text{rk}, \text{det}) : K_{0,X/k} \rightarrow \mathbf{Z}_{X/k} \times \text{Pic}_{X/k}$. If $\tau \in \{\text{Zar}, \text{ét}, \text{fppf}\}$, we let $K_{0,X/k,\tau}$ (resp. $\text{SK}_{0,X/k,\tau}$) be the sheafification of $K_{0,X/k}$ (resp. $\text{SK}_{0,X/k}$) for the corresponding (Zariski, étale, fppf) topology.

REMARK 2.2. – We do not know whether $K_{0,X/k,\text{ét}}$ and $K_{0,X/k,\text{fppf}}$ coincide.

2.2.2. *Functoriality.* – Let $f : X \rightarrow Y$ be a morphism of proper varieties over k . The pull-backs $(f_T)^*$ for $T \in (\text{Sch}/k)$ induce a natural transformation of functors $f^* : K_{0,Y/k} \rightarrow K_{0,X/k}$.

Similarly, if $f : X \rightarrow Y$ is a perfect (for instance lci) morphism of proper varieties over k , the push-forwards $(f_T)_*$ for $T \in (\text{Sch}/k)$ (which exist by §2.1.2 since f_T is perfect by [38, III, Corollaire 4.7.2]) induce a natural transformation of functors $f_* : K_{0,X/k} \rightarrow K_{0,Y/k}$, by the base change theorem [64, Theorem 3.10.3] (which can be applied since $f_T : X_T \rightarrow Y_T$ and $Y_U \rightarrow Y_T$ are Tor-independent for all morphisms $U \rightarrow T$ in (Sch/k)).

PROPOSITION 2.3. – *Let $f : X \rightarrow Y$ be a perfect birational morphism between proper integral varieties over k .*

- (i) *If Y is normal, then f_* restricts to a morphism $f_* : \text{SK}_{0,X/k} \rightarrow \text{SK}_{0,Y/k}$.*
- (ii) *If X and Y are regular, then $f_* \circ f^*$ is the identity of $K_{0,Y/k}$.*

Proof of (i). – Let $U \subseteq Y$ be the biggest open subset above which f is an isomorphism. Since Y is normal, the depth of $\mathcal{O}_{Y,y}$ is ≥ 2 for all $y \in Y \setminus U$.

Fix $T \in (\text{Sch}/k)$ and a class $x \in \text{SK}_0(X_T)$. Since $(\text{rk}(x), \text{det}(x))|_{U_T}$ is trivial, so is

$$(\text{rk}((f_T)_*x), \text{det}((f_T)_*x))|_{U_T} \in \mathbf{Z}(U_T) \times \text{Pic}(U_T).$$

To deduce the triviality of $(\text{rk}((f_T)_*x), \text{det}((f_T)_*x))$, it suffices to show the injectivity of the restriction morphisms $\mathbf{Z}(Y_T) \rightarrow \mathbf{Z}(U_T)$ and $\text{Pic}(Y_T) \rightarrow \text{Pic}(U_T)$.

The morphism $\mathbf{Z}(Y_T) \rightarrow \mathbf{Z}(U_T)$ is actually bijective by Remark 1.2 (ii) applied to the injection $U \hookrightarrow Y$. To show the injectivity of $\text{Pic}(Y_T) \rightarrow \text{Pic}(U_T)$, we can assume that T is noetherian by absolute noetherian approximation [80, Theorem C.9] and by the limit arguments of [36, §8.5]. It then suffices to combine [42, XI, Lemme 3.4] and [35, Proposition 6.3.1].

Proof of (ii). – Fix $T \in (\text{Sch}/k)$. Chatzistamatiou and Rülling [13, Theorem 1.1] have shown that the natural morphism $\mathcal{O}_Y \rightarrow Rf_*\mathcal{O}_X$ is a quasi-isomorphism. The base change theorem [64, Theorem 3.10.3] (which can be applied as Y_T is flat over Y) implies that the natural morphism $\mathcal{O}_{Y_T} \rightarrow R(f_T)_*\mathcal{O}_{X_T}$ is also a quasi-isomorphism. That $(f_T)_* \circ (f_T)^*$ is the identity of $K_0(Y_T)$ then follows from the projection formula (2.1). \square

2.2.3. *Curves.* – We can entirely compute $\mathbf{K}_{0,X/k,\tau}$ if X is a curve.

PROPOSITION 2.4. – *If $\tau \in \{\text{Zar}, \acute{\text{e}}\text{t}, \text{fppf}\}$ and if X is a projective variety of dimension ≤ 1 over k , then $(\text{rk}, \det) : \mathbf{K}_{0,X/k,\tau} \rightarrow \mathbf{Z}_{X/k} \times \text{Pic}_{X/k,\tau}$ is an isomorphism.*

Proof. – It suffices to prove the proposition for $\tau = \text{Zar}$. The commutation of \mathbf{K}_0 and Pic with directed inverse limits of qcqs schemes with affine transition maps (see [80, Proposition 3.20] and [36, §8.5]), applied to the system of affine neighborhoods of a point in a qcqs k -scheme, shows that it suffices to prove the bijectivity of $(\text{rk}, \det) : \mathbf{K}_0(X_T) \rightarrow \mathbf{Z}(X_T) \times \text{Pic}(X_T)$ for any local k -scheme T . By absolute noetherian approximation [80, Theorem C.9], we can write T as the limit of a directed inverse system $(T_i)_{i \in I}$ of noetherian k -schemes with affine transition maps. Replacing the T_i by their localizations at the images of the closed point of T , we may assume that they are local. A limit argument as above then shows that we may assume T to be noetherian. This case follows from Lemma 2.5 below applied to the connected components of X_T . \square

LEMMA 2.5. – *Let $\pi : Y \rightarrow T$ be a projective morphism with T local noetherian, Y non-empty and connected, and fibers of dimension ≤ 1 .*

Then the morphism $(\text{rk}, \det) : \mathbf{K}_0(Y) \rightarrow \mathbf{Z} \times \text{Pic}(Y)$ is bijective.

Proof. – The surjectivity of (rk, \det) is obvious and we prove its injectivity.

As explained in §2.1.1, $\mathbf{K}_0(Y)$ is generated by classes of vector bundles on Y . Lemma 2.6 below and induction on the rank of vector bundles show that it is even generated by classes of line bundles on Y . Let \mathcal{L} and \mathcal{M} be two line bundles on Y . Applying Lemma 2.6 twice with the same $l \gg 0$ and with the same very ample line bundle $\mathcal{O}_Y(1)$ on Y yields exact sequences $0 \rightarrow \mathcal{O}_Y(-l) \rightarrow \mathcal{L} \oplus \mathcal{M} \rightarrow \mathcal{F}_1 \rightarrow 0$ and $0 \rightarrow \mathcal{O}_Y(-l) \rightarrow \mathcal{O}_Y \oplus (\mathcal{L} \otimes \mathcal{M}) \rightarrow \mathcal{F}_2 \rightarrow 0$. Since the line bundles \mathcal{F}_1 and \mathcal{F}_2 have the same determinant, they are isomorphic, and we deduce the identity $[\mathcal{L}] + [\mathcal{M}] = [\mathcal{O}_Y] + [\mathcal{L} \otimes \mathcal{M}] \in \mathbf{K}_0(Y)$. This identity and the fact that $\mathbf{K}_0(Y)$ is generated by line bundles implies that $x = [\det(x)] + (\text{rk}(x) - 1)[\mathcal{O}_Y]$ for all $x \in \mathbf{K}_0(Y)$. This shows at once the required injectivity. \square

LEMMA 2.6. – *In the setting of Lemma 2.5, there exists a π -ample line bundle $\mathcal{O}_Y(1)$ on Y with the following property. For all vector bundles \mathcal{E} of rank ≥ 2 on Y and all $l \gg 0$, there exists a short exact sequence of vector bundles on Y of the form*

$$(2.6) \quad 0 \rightarrow \mathcal{O}_Y(-l) \rightarrow \mathcal{E} \rightarrow \mathcal{F} \rightarrow 0.$$

Proof. – Let $\mathcal{O}_Y(1)$ be a π -ample line bundle on Y , let t be the closed point of T , and let $A \subseteq Y_t$ be a finite subset meeting all the irreducible components of Y_t . If $m \gg 0$, then $H^0(Y_t, \mathcal{O}_{Y_t}(m)) \rightarrow H^0(A, \mathcal{O}_A(m))$ is surjective. As a consequence, after replacing $\mathcal{O}_Y(1)$ with $\mathcal{O}_Y(m)$, we may assume the existence of a section $\alpha \in H^0(Y_t, \mathcal{O}_{Y_t}(1))$ that vanishes at only finitely many points.

The same argument yields, for some $m \geq 0$, a section $\beta \in H^0(Y_t, \mathcal{E}_{Y_t}(m))$ that vanishes at only finitely many points. Let $B \subseteq Y_t$ be a finite subset meeting every irreducible component of Y_t and such that $\alpha\beta$ does not vanish at any point of B . Since $H^0(Y_t, \mathcal{E}_{Y_t}(n)) \rightarrow H^0(B, \mathcal{E}_B(n))$ is surjective for $n \gg 0$, we can choose $n \geq 0$ and a section $\gamma \in H^0(Y_t, \mathcal{E}_{Y_t}(n))$ such that $\alpha\beta$ and γ are linearly dependent at only finitely many points of Y_t . Let $C \subseteq Y_t$ be this finite set of points.

Choose any $l \geq \max(m, n)$ such that $H^0(Y_t, \mathcal{O}_{Y_t}(l-n)) \rightarrow H^0(C, \mathcal{O}_C(l-n))$ and $H^0(Y, \mathcal{E}(l)) \rightarrow H^0(Y_t, \mathcal{E}_{Y_t}(l))$ are surjective. (Such l exist by Serre vanishing [32, Théorème 2.2.1]; this is where we use the noetherianity of T .) Then there exists $\delta \in H^0(Y_t, \mathcal{O}_{Y_t}(l-n))$ such that $\tau := \alpha^{l-m}\beta + \gamma\delta \in H^0(Y_t, \mathcal{E}_{Y_t}(l))$ vanishes nowhere. Lift τ to a section $\sigma \in H^0(Y, \mathcal{E}(l))$. Since π is proper and T is local, σ does not vanish on Y , thus giving rise to a short exact sequence of the form (2.6). \square

REMARK 2.7. – If the residue field of T is infinite, the proof of Lemma 2.6 can be simplified as one can then apply [55, Corollary 3.6] on Y_t to construct τ .

2.2.4. *Projective bundles and blow-ups.* – If X is a proper variety over k and if \mathcal{E} is a vector bundle of rank r on X , the formula (2.2) induces an isomorphism of functors

$$(2.7) \quad \mathbf{K}_{0, X/k}^r \xrightarrow{\sim} \mathbf{K}_{0, \mathbf{P}_X \mathcal{E}/k}.$$

In view of the isomorphism $\mathrm{rk} : \mathbf{K}_{0, \mathrm{Spec}(k)/k, \mathrm{Zar}} \xrightarrow{\sim} \mathbf{Z}$ given by Proposition 2.4, we deduce that $\mathbf{K}_{0, \mathbf{P}_k^{r-1}/k, \mathrm{Zar}} \simeq \mathbf{Z}^r$ and that $([\mathcal{O}_{\mathbf{P}_k^{r-1}}(-j)])_{0 \leq j \leq r-1}$ forms a basis of the \mathbf{Z} -module $\mathbf{K}_{0, \mathbf{P}_k^{r-1}/k, \mathrm{Zar}}(k)$. The family $([\mathcal{O}_{\mathbf{P}_k^d}])_{0 \leq d \leq r-1}$ is another basis. For $r \geq 2$, identifying the morphism $(\mathrm{rk}, \det) : \mathbf{K}_{0, \mathbf{P}_k^{r-1}/k, \mathrm{Zar}} \rightarrow \mathbf{Z}_{\mathbf{P}_k^{r-1}/k} \times \mathrm{Pic}_{\mathbf{P}_k^{r-1}/k, \mathrm{Zar}} = \mathbf{Z}^2$ yields an isomorphism $\mathrm{SK}_{0, \mathbf{P}_k^{r-1}/k, \mathrm{Zar}} \simeq \mathbf{Z}^{r-2}$ and shows that $([\mathcal{O}_{\mathbf{P}_k^d}])_{0 \leq d \leq r-3}$ is a basis of the \mathbf{Z} -module $\mathrm{SK}_{0, \mathbf{P}_k^{r-1}/k, \mathrm{Zar}}(k)$.

Let X be a proper variety over k and $i : Y \rightarrow X$ a regular closed immersion of pure codimension $c \geq 1$. Define $p : X' \rightarrow X$ to be the blow-up of X along Y with exceptional divisor Y' , and $p' : Y' \rightarrow Y$ and $i' : Y' \rightarrow X'$ to be the natural morphisms. Then the formula (2.3) induces an isomorphism of functors

$$(2.8) \quad \mathbf{K}_{0, X/k} \times \mathbf{K}_{0, Y'/k}^{c-1} \xrightarrow{\sim} \mathbf{K}_{0, X'/k},$$

in view of the functorialities described in §2.2.2.

2.3. The functor $\mathrm{CH}_{X/k, \mathrm{fppf}}^2$

We introduce, for a smooth proper geometrically connected threefold X over k with geometrically trivial Chow group of zero-cycles, the functor $\mathrm{CH}_{X/k, \mathrm{fppf}}^2$. It will play for codimension 2 cycles the same role as the Picard functor $\mathrm{Pic}_{X/k, \mathrm{fppf}}$ does for codimension 1 cycles.

2.3.1. *The class of a point.* – We first exhibit a canonical class $\nu_X \in \mathbf{K}_{0, X/k, \mathrm{fppf}}(k)$.

PROPOSITION 2.8. – *Let X be a smooth proper geometrically connected variety over k whose degree map $\mathrm{deg} : \mathrm{CH}_0(X_{\bar{k}}) \rightarrow \mathbf{Z}$ is an isomorphism. Choose $\tau \in \{\mathrm{Zar}, \acute{\mathrm{e}}\mathrm{t}, \mathrm{fppf}\}$. Assume that $k = \bar{k}$ if $\tau = \mathrm{Zar}$ and that $k = k_{\mathfrak{p}}$ if $\tau = \acute{\mathrm{e}}\mathrm{t}$. There exists a unique $\nu_X \in \mathbf{K}_{0, X/k, \tau}(k)$ such that for all finite extensions k' of k and all coherent sheaves \mathcal{F} on $X_{k'}$ whose support has dimension 0, one has*

$$(2.9) \quad [\mathcal{F}] = h^0(X_{k'}, \mathcal{F}) \cdot \nu_X \in \mathbf{K}_{0, X/k, \tau}(k').$$

Proof. – Using [12, 2.2/13], choose a finite Galois extension l of k and a point $x \in X(l)$. Let n be the dimension of X . For all field extensions l' of l , the definition of the flat pull-back of a cycle [31, §1.5, §1.7] and the formula [38, X, (1.1.3)] show the commutativity of the natural diagram

$$(2.10) \quad \begin{array}{ccc} \mathrm{CH}_0(X_l) & \xrightarrow{\varphi^n} & F_0\mathrm{K}_0(X_l) \\ \downarrow & & \downarrow \\ \mathrm{CH}_0(X_{l'}) & \xrightarrow{\varphi^n} & F_0\mathrm{K}_0(X_{l'}), \end{array}$$

where the morphisms φ^n are defined in §2.1.5.

Assume first that $\tau = \text{fppf}$.

Let $\{x_1, \dots, x_m\}$ be the $\mathrm{Gal}(l/k)$ -orbit of x . Since $\mathrm{deg} : \mathrm{CH}_0(X_{\bar{k}}) \xrightarrow{\sim} \mathbf{Z}$ is an isomorphism, there exists a finite extension l' of l such that the x_i all have the same class in $\mathrm{CH}_0(X_{l'})$. Since $\mathrm{Spec}(l') \rightarrow \mathrm{Spec}(l)$ is an fppf covering, we deduce from (2.10) that $\varphi^n([x]) \in \mathrm{K}_{0,X/k,\tau}(l)$ is $\mathrm{Gal}(l/k)$ -invariant, hence descends to a class $\nu_X \in \mathrm{K}_{0,X/k,\tau}(k)$ since $\mathrm{Spec}(l) \rightarrow \mathrm{Spec}(k)$ is an étale covering. Applying (2.9) with $k' = l$ and $\mathcal{F} = \mathcal{O}_x$ shows that this is the only possible choice for ν_X and proves the uniqueness assertion of Proposition 2.8.

Let us now show that ν_X satisfies (2.9). Let k' be a finite extension of k and let \mathcal{F} be a coherent sheaf on $X_{k'}$ whose support has dimension 0. Let l' be a finite extension of k containing both k' and l , with the property that all the points in the support of $\mathcal{F}_{l'}$ have residue field l' , and are rationally equivalent to x . (Such an l' exists since $\mathrm{deg} : \mathrm{CH}_0(X_{\bar{k}}) \xrightarrow{\sim} \mathbf{Z}$.) The formula [38, X, (1.1.3)] and the commutativity of (2.10) show that $[\mathcal{F}_{l'}] = h^0(X_{l'}, \mathcal{F}_{l'}) \cdot \nu_X \in \mathrm{K}_{0,X/k,\tau}(l')$. Since $\mathrm{Spec}(l') \rightarrow \mathrm{Spec}(k')$ is an fppf covering, identity (2.9) follows.

If $\tau = \text{ét}$ (resp. $\tau = \text{Zar}$), all the fppf (resp. fppf or étale) coverings that appear above are étale (resp. Zariski) coverings, proving the proposition in these cases. \square

2.3.2. Codimension 2 cycles on a threefold. – Let us fix in §2.3.2 a smooth proper geometrically connected threefold X over k whose degree map $\mathrm{deg} : \mathrm{CH}_0(X_{\Omega}) \rightarrow \mathbf{Z}$ is an isomorphism for all algebraically closed field extensions $k \subseteq \Omega$ (an assumption that is satisfied if X is \bar{k} -rational).

Choose $\tau \in \{\text{Zar}, \text{ét}, \text{fppf}\}$, and assume that $k = \bar{k}$ if $\tau = \text{Zar}$ and that $k = k_{\mathfrak{p}}$ if $\tau = \text{ét}$, so that $\mathrm{Spec}(l) \rightarrow \mathrm{Spec}(k)$ is a τ -covering for any finite extension l of k . Let $\nu_X \in \mathrm{K}_{0,X/k,\tau}(k)$ be the class defined in Proposition 2.8. If $x \in X(l)$ for some finite extension l of k , then $(\mathrm{rk}, \mathrm{det})([\mathcal{O}_x]) = (0, \mathcal{O}_X)$ (see §2.1.5). In view of (2.9), one therefore has $\nu_X \in \mathrm{SK}_{0,X/k,\tau}(k) \subseteq \mathrm{SK}_{0,X/k,\tau}(l)$.

We still denote by ν_X the morphism of τ -sheaves $\nu_X : \mathbf{Z} \rightarrow \mathrm{SK}_{0,X/k,\tau}$ such that $\nu_X(1) = \nu_X$. We view ν_X as a morphism of presheaves of abelian groups.

DEFINITION 2.9. – We let $\mathrm{CH}_{X/k,\tau}^2 : (\mathrm{Sch}/k)^{\mathrm{op}} \rightarrow (\mathrm{Ab})$ be the (presheaf) cokernel of $\nu_X : \mathbf{Z} \rightarrow \mathrm{SK}_{0,X/k,\tau}$. When $\mathrm{CH}_{X/k,\mathrm{fppf}}^2$ is representable, we let $\mathbf{CH}_{X/k}^2$ be the group scheme over k that represents it.

REMARK 2.10. – Let k'/k be a field extension. There is an obvious identification $\mathbf{K}_{0, X_{k'}/k'}(T) = \mathbf{K}_{0, X/k}(T)$ for all $T \in (\text{Sch}/k')$. We thus obtain natural isomorphisms $\mathbf{K}_{0, X_{k'}/k', \tau}(T) = \mathbf{K}_{0, X/k, \tau}(T)$, $\mathbf{SK}_{0, X_{k'}/k', \tau}(T) = \mathbf{SK}_{0, X/k, \tau}(T)$ and $\mathbf{CH}_{X_{k'}/k', \tau}^2(T) = \mathbf{CH}_{X/k, \tau}^2(T)$ for all $T \in (\text{Sch}/k')$ and $\tau \in \{\text{Zar}, \acute{\text{e}}\text{t}, \text{fppf}\}$. In particular, if $\mathbf{CH}_{X/k, \text{fppf}}^2$ is representable, so is $\mathbf{CH}_{X_{k'}/k', \text{fppf}}^2$ and $\mathbf{CH}_{X_{k'}/k'}^2 = (\mathbf{CH}_{X/k}^2)_{k'}$.

The following proposition justifies these definitions.

PROPOSITION 2.11. – *Associating with the class $[Z] \in \mathbf{CH}^2(X_{\bar{k}})$ of a codimension 2 integral closed subvariety $Z \subset X_{\bar{k}}$ the class $[\mathcal{O}_Z] \in \mathbf{K}_0(X_{\bar{k}})$ of its structure sheaf induces a Γ_k -equivariant isomorphism*

$$(2.11) \quad \mathbf{CH}^2(X_{\bar{k}}) \xrightarrow{\sim} \mathbf{CH}_{X_{\bar{k}}/\bar{k}, \text{Zar}}^2(\bar{k}) = \mathbf{CH}_{X/k, \text{Zar}}^2(\bar{k}).$$

Proof. – In view of (2.9), one has a natural isomorphism

$$(2.12) \quad \mathbf{SK}_0(X_{\bar{k}})/F_0\mathbf{K}_0(X_{\bar{k}}) \xrightarrow{\sim} \mathbf{CH}_{X_{\bar{k}}/\bar{k}, \text{Zar}}^2(\bar{k}).$$

Precomposing (2.12) with $\varphi^2 : \mathbf{CH}^2(X_{\bar{k}}) \xrightarrow{\sim} \text{Gr}_F^2 \mathbf{K}_0(X_{\bar{k}}) = \mathbf{SK}_0(X_{\bar{k}})/F_0\mathbf{K}_0(X_{\bar{k}})$ (see §2.1.5) yields the isomorphism (2.11). It is Γ_k -equivariant by construction. \square

The next lemma will be used in the proof of Theorem 3.1 (iv). For the definition of $\alpha|_{X_t} \in \mathbf{CH}^2(X_t)$ in its statement, see [31, Example 5.2.1].

LEMMA 2.12. – *Let T be a smooth connected variety over \bar{k} .*

- (i) *For all $\alpha \in \mathbf{CH}^2(X_T)$, there exists a class $\beta \in \mathbf{SK}_0(X_T)$ with the property that for all $t \in T(\bar{k})$, the image of $\alpha|_{X_t}$ by (2.11) is induced by $\beta|_{X_t}$.*
- (ii) *For all $\beta \in \mathbf{SK}_0(X_T)$, there exists a class $\alpha \in \mathbf{CH}^2(X_T)$ with the property that for all $t \in T(\bar{k})$, the image of $\alpha|_{X_t}$ by (2.11) is induced by $\beta|_{X_t}$.*

Proof of (i). – We can assume that α is the class of an integral subvariety $Z \subset X_T$ of codimension 2. Define $\beta := [\mathcal{O}_Z] \in \mathbf{SK}_0(X_T)$. By the Riemann-Roch theorem without denominators, one has $\alpha = -c_2(\beta)$ (see §2.1.5). For $t \in T(\bar{k})$, one has $\alpha|_{X_t} = -c_2(\beta|_{X_t})$. Applying the Riemann-Roch theorem without denominators again shows that the image of $\alpha|_{X_t}$ by (2.11) is the class induced by $\beta|_{X_t}$.

Proof of (ii). – Define $\alpha = -c_2(\beta) \in \mathbf{CH}^2(X_T)$ and argue as in (i). \square

3. Geometrically rational threefolds

In Section 3, we prove the representability of the functor $\mathbf{CH}_{X/k, \text{fppf}}^2$ defined in §2.3.2 if X is a smooth projective \bar{k} -rational threefold, and study the group scheme $\mathbf{CH}_{X/k}^2$ that represents it.

3.1. Main statement

Our goal is the following theorem.

THEOREM 3.1. – *Let X be a smooth projective \bar{k} -rational threefold over k . Then:*

- (i) $\mathrm{CH}_{X/k, \mathrm{fppf}}^2$ is represented by a smooth group scheme $\mathbf{CH}_{X/k}^2$ over k .
- (ii) $(\mathbf{CH}_{X/k}^2)^0$ is an abelian variety over k .
- (iii) $\mathrm{CH}_{X_{k_p}/k_p, \mathrm{fppf}}^2 = \mathrm{CH}_{X_{k_p}/k_p, \acute{\mathrm{e}}\mathrm{t}}^2$ and $\mathrm{CH}_{X_{\bar{k}}/\bar{k}, \mathrm{fppf}}^2 = \mathrm{CH}_{X_{\bar{k}}/\bar{k}, \mathrm{Zar}}^2$.
- (iv) The Γ_k -equivariant isomorphism

$$(3.1) \quad \psi_X^2 : \mathrm{CH}^2(X_{\bar{k}}) \xrightarrow{\sim} \mathbf{CH}_{X/k}^2(\bar{k})$$
 obtained by combining (iii) and (2.11) restricts to a Γ_k -equivariant bijective regular homomorphism (in the sense of §1.6)

$$(3.2) \quad \psi_X^2 : \mathrm{CH}^2(X_{\bar{k}})_{\mathrm{alg}} \xrightarrow{\sim} (\mathbf{CH}_{X/k}^2)^0(\bar{k}).$$
- (v) The étale group scheme $\mathbf{CH}_{X/k}^2 / (\mathbf{CH}_{X/k}^2)^0$ over k is associated with the Γ_k -module $\mathrm{NS}^2(X_{\bar{k}})$, which as a \mathbf{Z} -module is free of finite rank.
- (vi) The isomorphism (3.2) induces an isomorphism $\mathrm{Ab}^2(X_{k_p}) \xrightarrow{\sim} (\mathbf{CH}_{X_{k_p}/k_p}^2)^0$ (where $\mathrm{Ab}^2(X_{k_p})$ denotes Murre's intermediate Jacobian, introduced in §1.6).

We endow $\mathbf{CH}_{X/k}^2$ with the principal polarization (in the sense of §1.2) induced by the principal polarization θ_X of $\mathrm{Ab}^2(X_{k_p})$ (see §1.6).

- (vii) If X is k -rational, there exists a smooth projective curve B over k such that $\mathbf{CH}_{X/k}^2$ is a principally polarized direct factor of $\mathbf{Pic}_{B/k}$ (in the sense of §1.2).

The proof of Theorem 3.1 is given in §3.2. Theorem 3.1 is complemented in §3.3 by a computation of $\mathbf{CH}_{X/k}^2$ for varieties constructed as blow-ups, and in §3.4 by an analysis of the obstructions to k -rationality arising from Theorem 3.1 (vii).

REMARKS 3.2. – (i) Let X be a smooth projective variety over k . As recalled in §1.6, Achter, Casalaina-Martin and Vial have endowed $\mathrm{Ab}^2(X_{\bar{k}})$ with a natural k_p -structure. If X is moreover a \bar{k} -rational threefold, Theorem 3.1 (vi) further endows $\mathrm{Ab}^2(X_{\bar{k}})$ with a natural k -structure $(\mathbf{CH}_{X/k}^2)^0$. Trying to descend $\mathrm{Ab}^2(X_{\bar{k}})$ to k under more general hypotheses gives an incentive to define $\mathrm{CH}_{X/k, \mathrm{fppf}}^2$ and to prove its representability in a greater generality.

(ii) For instance it would be nice to define and study a functor $\mathrm{CH}_{X/k, \mathrm{fppf}}^2$ for all smooth proper varieties X over k such that $\mathrm{CH}_0(X)_{\mathbf{Q}}$ is supported in dimension 1 in the sense of Bloch and Srinivas (see [7, Definition 2.1]). If a good enough theory of motivic cohomology $H_{\mathcal{M}}^*$ over a field of characteristic $p > 0$ were available without having to invert p in the coefficients, a natural choice would be the fppf sheafification of the functor $T \mapsto H_{\mathcal{M}}^4(X_T, \mathbf{Z}(2))$. One could also consider the fppf sheafification of the functor $T \mapsto H^2(X_T, \mathcal{K}_2)$, where \mathcal{K}_2 denotes Quillen's K -theory sheaf [9], or of the functor $T \mapsto A^2(X_T)$, where A^2 denotes Fulton's cohomological Chow group [31, Definition 17.3].

(iii) Even with our definition of $\mathrm{CH}_{X/k, \mathrm{fppf}}^2$, it would be interesting to show the representability of $\mathrm{CH}_{X/k, \mathrm{fppf}}^2$ under the hypothesis, weaker than \bar{k} -rationality, that X is a smooth,

proper and geometrically connected threefold over k such that $\text{deg} : \text{CH}_0(X_\Omega) \xrightarrow{\sim} \mathbf{Z}$ is an isomorphism for all algebraically closed field extensions $k \subseteq \Omega$. We note, however, that it is the proof of representability that we give here, and which is specific to \bar{k} -rational threefolds, that yields the crucial Theorem 3.1 (vii) and thus provides obstructions to k -rationality.

(iv) We still denote by θ_X the principal polarization of $\mathbf{CH}_{X/k}^2$ induced by that of $\text{Ab}^2(X_{k_p})$, as in Theorem 3.1 (vi). For the sake of completeness, we extract a characterization of θ_X from the definition of the isomorphism (3.2) and from [7, Property 2.4, Corollary 2.8]. Let ℓ be a prime number invertible in k . Consider the diagram

$$(\mathbf{CH}_{X/k}^2)^0(\bar{k})\{\ell\} \xleftarrow[\sim]{\psi_{\bar{X}}^2} \text{CH}^2(X_{\bar{k}})_{\text{alg}}\{\ell\} \xrightarrow{\lambda^2} H^3(X_{\bar{k}}, \mathbf{Z}_\ell(2)) \otimes \mathbf{Q}_\ell/\mathbf{Z}_\ell,$$

where λ^2 is Bloch's Abel-Jacobi map (see [10, §2], [7, (2.3)]) and $\psi_{\bar{X}}^2$ is the map (3.2). Then $c_1(\theta_X) \in H^2((\mathbf{CH}_{X/k}^2)^0_{\bar{k}}, \mathbf{Z}_\ell(1)) = (\wedge^2 H^1((\mathbf{CH}_{X/k}^2)^0_{\bar{k}}, \mathbf{Z}_\ell))(1)$ corresponds, via the identification

$$H^1((\mathbf{CH}_{X/k}^2)^0_{\bar{k}}, \mathbf{Z}_\ell)^\vee \xrightarrow{T_\ell(\lambda^2 \circ (\psi_{\bar{X}}^2)^{-1})} H^3(X_{\bar{k}}, \mathbf{Z}_\ell(2))/(\text{torsion})$$

(in which $^\vee$ stands for $\text{Hom}(-, \mathbf{Z}_\ell)$), to the opposite of the cup product pairing

$$\bigwedge^2 H^3(X_{\bar{k}}, \mathbf{Z}_\ell(2)) \rightarrow H^6(X_{\bar{k}}, \mathbf{Z}_\ell(4)) \xrightarrow{\sim} \mathbf{Z}_\ell(1).$$

3.2. Proof of Theorem 3.1

3.2.1. *Resolution of indeterminacies.* – Our main tool is a resolution of indeterminacies result that is due to Abhyankar [1] if k is perfect, and to Cossart and Piltant [22] in general.

PROPOSITION 3.3. – *Let X and Y be smooth projective threefolds over k . Let $f : Y \dashrightarrow X$ be a birational map. Then there exists a diagram*

$$(3.3) \quad X \xleftarrow{h} X' = Y_{N+1} \rightarrow \cdots \rightarrow Y_{j+1} \xrightarrow{p_j} Y_j \rightarrow \cdots \rightarrow Y_1 = Y$$

of regular projective varieties over k such that $f = h \circ p_N^{-1} \circ \cdots \circ p_1^{-1}$, where p_j is the blow-up of an integral regular closed subscheme $Z_j \subset Y_j$ of codimension c_j and where h is projective and birational.

Proof. – This follows from [22], as explained in [7, Proposition 2.11]. The standing assumption that k is perfect in [7] is irrelevant if one really uses [22, Proposition 4.2] (or [21, Theorem 5.9]) instead of [1, (9.1.4)] in the proof of [7, Proposition 2.11]. \square

REMARK 3.4. – If k is not perfect, the subschemes $Z_j \subset Y_j$ may not be smooth over k .

3.2.2. *Representability if X is k -rational.* – In §3.2.2, we fix $\tau \in \{\text{Zar}, \text{ét}, \text{fppf}\}$ and assume that $k = \bar{k}$ if $\tau = \text{Zar}$ and that $k = k_p$ if $\tau = \text{ét}$. We also let X be a smooth projective k -rational threefold. By Proposition 3.3, there exists a diagram (3.3) with $Y = \mathbf{P}_k^3$. Remark 1.2 (ii), Corollary 1.5 and the isomorphism (2.8) then give canonical decompositions

$$(3.4) \quad \mathbf{Z}_{X'/k} \xleftarrow{\sim} \mathbf{Z}_{\mathbf{P}_k^3/k},$$

$$(3.5) \quad \text{Pic}_{X'/k} \xleftarrow{\sim} \text{Pic}_{\mathbf{P}_k^3/k} \times \prod_{c_j \geq 2} \mathbf{Z}_{Z_j/k},$$

$$(3.6) \quad \mathbf{K}_{0,X'/k} \xleftarrow{\sim} \mathbf{K}_{0,\mathbf{P}_k^3/k} \times \prod_{c_j=2} \mathbf{K}_{0,Z_j/k} \times \prod_{c_j=3} (\mathbf{K}_{0,Z_j/k})^2.$$

Identifying $(\text{rk}, \det) : \mathbf{K}_{0,X'/k} \rightarrow \mathbf{Z}_{X'/k} \times \text{Pic}_{X'/k}$ in terms of these decompositions and using the isomorphisms $(\text{rk}, \det) : \mathbf{K}_{0,Z_j/k,\tau} \xrightarrow{\sim} \mathbf{Z}_{Z_j/k} \times \text{Pic}_{Z_j/k,\tau}$ given by Proposition 2.4 yields an isomorphism

$$(3.7) \quad \text{SK}_{0,X'/k,\tau} \xleftarrow{\sim} \text{SK}_{0,\mathbf{P}_k^3/k,\tau} \times \prod_{c_j=2} \text{Pic}_{Z_j/k,\tau} \times \prod_{c_j=3} \mathbf{Z}_{Z_j/k}.$$

The morphisms p_j are lci by [79, §1.2], hence so is the structural morphism $X' \rightarrow \text{Spec}(k)$ by [38, VIII, Proposition 1.5]. Any closed embedding $X' \hookrightarrow \mathbf{P}_X^N$ of the X -scheme X' is a regular immersion by [38, VIII, Proposition 1.2], which shows that h is lci, hence perfect [38, VIII, Proposition 1.7]. The functors $h^* : \mathbf{K}_{0,X/k} \rightarrow \mathbf{K}_{0,X'/k}$ and $h_* : \mathbf{K}_{0,X'/k} \rightarrow \mathbf{K}_{0,X/k}$ satisfy $h_* \circ h^* = \text{Id}$ by Proposition 2.3 (ii). Since they restrict to $h^* : \text{SK}_{0,X/k} \rightarrow \text{SK}_{0,X'/k}$ and to $h_* : \text{SK}_{0,X'/k} \rightarrow \text{SK}_{0,X/k}$ (see Proposition 2.3 (i)), we deduce a natural decomposition

$$(3.8) \quad \text{SK}_{0,X/k} \times \text{Ker}(h_* : \text{SK}_{0,X'/k} \rightarrow \text{SK}_{0,X/k}) \xrightarrow{\sim} \text{SK}_{0,X'/k}.$$

The three summands of the right-hand side of (3.7) are represented by group schemes locally of finite type over k , respectively by §2.2.4, by §1.4 and by Proposition 1.1. It follows that $\text{SK}_{0,X'/k,\tau}$ is represented by a group scheme locally of finite type over k . So is $\text{SK}_{0,X/k,\tau}$, by (3.8) and Lemma 1.17.

Let $x' \in X'(\bar{k})$ be a general point, and let $x \in X(\bar{k})$ and $y \in \mathbf{P}^3(\bar{k})$ be its images by h and by $p_1 \circ \dots \circ p_N$. Then $h^*[\mathcal{O}_x] = [\mathcal{O}_{x'}] = p_N^* \circ \dots \circ p_1^*[\mathcal{O}_y] \in \text{SK}_{0,X'/k,\tau}(\bar{k})$. Consequently, $h^* \circ \nu_X : \mathbf{Z} \rightarrow \text{SK}_{0,X'/k,\tau}$ and $p_N^* \circ \dots \circ p_1^* \circ \nu_{\mathbf{P}_k^3} : \mathbf{Z} \rightarrow \text{SK}_{0,X'/k,\tau}$ both send $1 \in \mathbf{Z}(\bar{k})$ to $[\mathcal{O}_{x'}] \in \text{SK}_{0,X'/k,\tau}(\bar{k})$. For $n \neq 0$, the class $\nu_{\mathbf{P}_k^3}(n)$ does not belong to the identity component of $\text{SK}_{0,\mathbf{P}_k^3/k,\tau}$, by §2.2.4. We deduce from the above and from (3.7) that $n[\mathcal{O}_{x'}] \notin (\text{SK}_{0,X'/k,\tau})^0(\bar{k})$, hence that $\nu_X(n) \notin (\text{SK}_{0,X/k,\tau})^0(\bar{k})$. Lemma 1.18 then shows that $\text{CH}_{X'/k,\tau}^2$ is represented by a group scheme locally of finite type over k .

Applying the above with $\tau = \text{fppf}$ and combining (3.7), (3.8) and the equality $h^* \circ \nu_X = p_N^* \circ \dots \circ p_1^* \circ \nu_{\mathbf{P}_k^3}$ yields isomorphisms

$$(3.9) \quad \mathbf{CH}_{X'/k}^2 \times G \xrightarrow{\sim} \mathbf{CH}_{X'/k}^2 \xleftarrow{\sim} \mathbf{CH}_{\mathbf{P}_k^3/k}^2 \times \prod_{c_j=2} \text{Pic}_{Z_j/k} \times \prod_{c_j=3} \mathbf{Z}_{Z_j/k}$$

of smooth group schemes over k , where G denotes the group scheme representing the fppf sheafification of $\text{Ker}(h_* : \text{SK}_{0,X'/k} \rightarrow \text{SK}_{0,X/k})$ (whose representability follows from

Lemma 1.17) and where $\mathbf{CH}_{X'/k}^2$ denotes the group scheme representing the presheaf cokernel $\text{Coker}(h^* \circ \nu_X : \mathbf{Z} \rightarrow \mathbf{SK}_{0,X'/k,\text{fppf}})$ (whose representability follows from Lemma 1.18).

3.2.3. *Representability if X is \bar{k} -rational.* – In §3.2.3, we prove Theorem 3.1 (i)–(iii) for a smooth projective \bar{k} -rational threefold X over k .

Choose $\tau \in \{\text{Zar}, \text{ét}, \text{fppf}\}$ and assume that $k = \bar{k}$ if $\tau = \text{Zar}$ and that $k = k_p$ if $\tau = \text{ét}$. Let l be a finite extension of k such that X is l -rational. Then $\mathbf{SK}_{0,X_l/l,\tau}$ is represented by a group scheme locally of finite type over l , by the arguments of §3.2.2 applied to the l -variety X_l . By Lemma 1.19, it follows that $\mathbf{SK}_{0,X/k,\tau}$ is represented by a group scheme locally of finite type over k . As explained in §3.2.2, the morphism $\nu_X : \mathbf{Z} \rightarrow \mathbf{SK}_{0,X/k,\tau}$ defined in §2.3.2 has the property that $\nu_X(n) \notin (\mathbf{SK}_{0,X/k,\tau})^0(\bar{k})$ for all $n \neq 0$. It now follows from Lemma 1.18 that $\mathbf{CH}_{X/k,\tau}^2$ is represented by a group scheme locally of finite type over k .

Proof of Theorem 3.1 (i)–(iii). – Applying the above argument to the k -variety X with $\tau = \text{fppf}$, to the k_p -variety X_{k_p} with $\tau = \text{ét}$, and to the \bar{k} -variety $X_{\bar{k}}$ with $\tau = \text{Zar}$ shows that the three functors $\mathbf{CH}_{X/k,\text{fppf}}^2$, $\mathbf{CH}_{X_{k_p}/k_p,\text{ét}}^2$ and $\mathbf{CH}_{X_{\bar{k}}/\bar{k},\text{Zar}}^2$ are represented by group schemes locally of finite type over k , over k_p and over \bar{k} , respectively. In particular, the latter two are sheaves for the fppf topology, which proves Theorem 3.1 (iii). In addition, the arguments of §3.2.2 applied to the \bar{k} -variety $X_{\bar{k}}$ show that $(\mathbf{CH}_{X_{\bar{k}}/\bar{k}}^2)^0$ is a direct factor of a product of Jacobians of smooth projective curves over \bar{k} (see (3.9)), hence is an abelian variety; in particular, it is smooth. As $((\mathbf{CH}_{X/k}^2)^0)_{\bar{k}} = (\mathbf{CH}_{X_{\bar{k}}/\bar{k}}^2)^0$, Theorem 3.1 (i)–(ii) follows. \square

3.2.4. *Relation with Murre's work.* – We now prove Theorem 3.1 (iv)–(vi).

Proof of Theorem 3.1 (iv). – That $\psi_X^2(\mathbf{CH}^2(X_{\bar{k}})_{\text{alg}}) \subseteq ((\mathbf{CH}_{X/k}^2)^0(\bar{k}))$, and that the resulting morphism $\psi_X^2 : \mathbf{CH}^2(X_{\bar{k}})_{\text{alg}} \rightarrow (\mathbf{CH}_{X/k}^2)^0(\bar{k})$ is a regular homomorphism follow at once from Lemma 2.12 (i). It remains to show that (3.2) is surjective.

Since $\mathbf{CH}_{X_{\bar{k}}/\bar{k}}^2$ represents $\mathbf{CH}_{X_{\bar{k}}/\bar{k},\text{Zar}}^2$ by Theorem 3.1 (iii), we can choose a connected Zariski neighborhood T of the identity in $(\mathbf{CH}_{X_{\bar{k}}/\bar{k}}^2)^0$ and a class $\beta \in \mathbf{SK}_0(X_T)$ inducing the natural inclusion $T \rightarrow \mathbf{CH}_{X_{\bar{k}}/\bar{k}}^2$. Lemma 2.12 (ii) then implies that $T(\bar{k})$ is contained in the image of (3.2). Since $(\mathbf{CH}_{X/k}^2)^0(\bar{k})$ is generated by $T(\bar{k})$ as a group, we have proved the surjectivity of (3.2). \square

Proof of Theorem 3.1 (v). – That the étale k -group scheme $(\mathbf{CH}_{X/k}^2)/(\mathbf{CH}_{X/k}^2)^0$ corresponds to the Γ_k -module $\text{NS}^2(X_{\bar{k}})$ follows at once from (3.1) and (3.2).

Applying the discussion of §3.2.2, and more precisely identity (3.9), to the \bar{k} -variety $X_{\bar{k}}$, shows, in view of the isomorphism $\mathbf{CH}_{\mathbf{P}_{\bar{k}}^3/\bar{k}}^2 \simeq \mathbf{Z}$ (see §2.2.4), that $\text{NS}^2(X_{\bar{k}})$ is a free \mathbf{Z} -module of finite rank, being a direct factor of such a module. \square

Proof of Theorem 3.1 (vi). – The regular homomorphism (3.2) induces a morphism $\iota_{X_{\bar{k}}} : \text{Ab}^2(X_{\bar{k}}) \rightarrow (\mathbf{CH}_{X_{\bar{k}}/\bar{k}}^2)^0$. Since (3.2) is Γ_k -equivariant, and in view of the definition of $\text{Ab}^2(X_{k_p})$ recalled in §1.6, this morphism descends by Galois descent to a morphism $\iota_{X_{k_p}} : \text{Ab}^2(X_{k_p}) \rightarrow (\mathbf{CH}_{X_{k_p}/k_p}^2)^0$ of abelian varieties over k_p . To prove that $\iota_{X_{k_p}}$ is an

isomorphism, it suffices to prove that $\iota_{X_{\bar{k}}}$ is an isomorphism. From now on, we may thus assume that $k = \bar{k}$.

By Proposition 3.3, there exists a diagram (3.3). Since $k = \bar{k}$, all the varieties Z_j and Y_j that appear in it are smooth over k . Consider the diagram

$$(3.10) \quad \begin{array}{ccc} \mathrm{Ab}^2(\mathbf{P}_k^3) \times \prod_{c_j=2} \mathbf{Pic}_{Z_j/k}^0 & \xrightarrow{\sim} & \mathrm{Ab}^2(X') \\ \downarrow (\iota_{\mathbf{P}_k^3}, \mathrm{Id}) & & \downarrow \iota_{X'} \\ (\mathbf{CH}_{\mathbf{P}_k^3/k}^2)^0 \times \prod_{c_j=2} \mathbf{Pic}_{Z_j/k}^0 & \xrightarrow{\sim} & (\mathbf{CH}_{X'/k}^2)^0, \end{array}$$

where the lower horizontal isomorphism is induced by (3.7) and the upper horizontal isomorphism is the one provided by [7, Lemma 2.10]. Since $\mathrm{CH}^2(\mathbf{P}_k^3)_{\mathrm{alg}} = 0$, one has $\mathrm{Ab}^2(\mathbf{P}_k^3) = (\mathbf{CH}_{\mathbf{P}_k^3/k}^2)^0 = 0$ and the left vertical arrow of (3.10) is an isomorphism. We claim that (3.10) commutes. Since $k = \bar{k}$, it suffices to verify that it commutes at the level of k -points, which follows from unwinding the definitions and making use of Lemma 2.1. A glance at (3.10) now shows that $\iota_{X'}$ is an isomorphism.

Now, consider the diagram

$$(3.11) \quad \begin{array}{ccccc} \mathrm{Ab}^2(X) & \xrightarrow{h^+} & \mathrm{Ab}^2(X') & \xrightarrow{h_+} & \mathrm{Ab}^2(X) \\ \downarrow \iota_X & & \wr \downarrow \iota_{X'} & & \downarrow \iota_X \\ (\mathbf{CH}_{X/k}^2)^0 & \xrightarrow{h^*} & (\mathbf{CH}_{X'/k}^2)^0 & \xrightarrow{h_*} & (\mathbf{CH}_{X/k}^2)^0, \end{array}$$

whose lower horizontal arrows are induced by (3.8) and hence satisfy $h_* \circ h^* = \mathrm{Id}$, and whose upper horizontal arrows are given by the functoriality of Murre’s intermediate Jacobians (see [7, §2.2.1]) and satisfy $h_+ \circ h^+ = \mathrm{Id}$ as a consequence of the identity $h_* \circ h^* = \mathrm{Id} : \mathrm{CH}^2(X) \rightarrow \mathrm{CH}^2(X)$ stemming from the projection formula [31, Proposition 8.3(c)]. To show that (3.11) commutes, it suffices to check that it commutes at the level of k -points, since $k = \bar{k}$. This follows from Lemma 2.1 for the left-hand square, and from the fact that the morphisms φ^c considered in §2.1.5 are compatible with proper push-forwards [31, Example 15.1.5] for the right-hand square. A diagram chase in (3.11) shows that ι_X is an isomorphism since $\iota_{X'}$ is one, which concludes the proof. \square

3.2.5. *Further analysis of k -rational varieties.* – We resume the discussion of §3.2.2 with $\tau = \mathrm{fppf}$, and keep the notation introduced there. Since $\mathbf{CH}_{\mathbf{P}_k^3/k}^2 \simeq \mathbf{Z}$ by §2.2.4, identity (3.9) reads:

$$(3.12) \quad \mathbf{CH}_{X/k}^2 \times G \xrightarrow{\sim} \mathbf{CH}_{X'/k}^2 \xleftarrow{\sim} \mathbf{Z} \times \prod_{c_j=2} \mathbf{Pic}_{Z_j/k} \times \prod_{c_j=3} \mathbf{Z}_{Z_j/k}.$$

The identity component of the right-hand side is isomorphic to $\prod_{c_j=2} \mathbf{Pic}_{Z_j/k}^0$, hence it carries a natural principal polarization (see §1.4.2). Via (3.12), we thus obtain a principal polarization on $\mathbf{CH}_{X'/k}^2$ in the sense of §1.2.

PROPOSITION 3.5. – *The isomorphism (3.12) realizes $\mathbf{CH}_{X'/k}^2$ and G as principally polarized direct factors, in the sense of §1.2, of $\mathbf{CH}_{X'/k}^2$, and the induced polarization on $\mathbf{CH}_{X'/k}^2$ coincides with the one defined in Theorem 3.1 (vi).*

Proof. – We fix once and for all a prime number ℓ invertible in k and start with a few recollections about (Borel-Moore) ℓ -adic étale homology. If V is a variety over \bar{k} , the i -th étale homology group of V with coefficients in $\mathbf{Q}_\ell(j)$ is defined by $H_i(V, \mathbf{Q}_\ell(j)) = H^{-i}(V, R\varepsilon^! \mathbf{Q}_\ell(j))$, where $\varepsilon : V \rightarrow \text{Spec}(\bar{k})$ denotes the structural morphism (see [63, Définition 2]). This group is covariantly functorial with respect to proper morphisms (*loc. cit.*, §4) and comes with a cap product operation $H^s(V, \mathbf{Q}_\ell(t)) \times H_i(V, \mathbf{Q}_\ell(j)) \rightarrow H_{i-s}(V, \mathbf{Q}_\ell(j+t))$ (*loc. cit.*, p. VIII-09), and with a cycle class map $\text{cl} : \text{CH}_i(V) \rightarrow H_{2i}(V, \mathbf{Q}_\ell(-i))$ (*loc. cit.*, §6), for any i, j, s, t . If V is of pure dimension n , we denote by $[V]$ the fundamental cycle of V (see [31, §1.5]), so that $\text{cl}([V]) \in H_{2n}(V, \mathbf{Q}_\ell(-n))$. For all i, j , the map

$$(3.13) \quad \kappa_V^{\text{ét}} : H^{2n-i}(V, \mathbf{Q}_\ell(j+n)) \rightarrow H_i(V, \mathbf{Q}_\ell(j))$$

defined by $\kappa_V^{\text{ét}}(x) = x \cap \text{cl}([V])$ is an isomorphism if in addition V is smooth (see [63, Prop. 3.2]). Thus, given a proper morphism $f : V' \rightarrow V$ from a variety V' of pure dimension n' to a smooth variety V of pure dimension n over \bar{k} , one can define a push-forward in étale cohomology

$$(3.14) \quad f_* : H^s(V', \mathbf{Q}_\ell(t)) \rightarrow H^{s+2n-2n'}(V, \mathbf{Q}_\ell(t+n-n'))$$

as the composition $(\kappa_V^{\text{ét}})^{-1} \circ f_* \circ \kappa_{V'}^{\text{ét}}$. When V is a proper variety over \bar{k} of pure odd dimension n , we will speak of the “cup product pairing on $H^n(V, \mathbf{Q}_\ell((n+1)/2))$ ” to refer to the pairing $H^n(V, \mathbf{Q}_\ell((n+1)/2)) \times H^n(V, \mathbf{Q}_\ell((n+1)/2)) \rightarrow \mathbf{Q}_\ell(1)$ induced by the cup product and by the push-forward map $H^{2n}(V, \mathbf{Q}_\ell(n+1)) \rightarrow \mathbf{Q}_\ell(1)$ along the structural morphism $V \rightarrow \text{Spec}(\bar{k})$ (see (3.14)).

LEMMA 3.6. – *Let $f : D \rightarrow V$ be a morphism between projective varieties over \bar{k} , where D has pure dimension d and V has pure dimension $d+1$, for some integer d . Suppose that V is smooth.*

Let $\lambda^1 : \text{Pic}(D)\{\ell\} \xrightarrow{\sim} H^1(D, \mathbf{Q}_\ell/\mathbf{Z}_\ell(1))$ and $\lambda^2 : \text{CH}^2(V)\{\ell\} \rightarrow H^3(V, \mathbf{Q}_\ell/\mathbf{Z}_\ell(2))$ respectively denote the Kummer isomorphism and Bloch’s ℓ -adic Abel-Jacobi map. Then the diagram

$$(3.15) \quad \begin{array}{ccccc} V_\ell(F^2 \mathbf{K}_0(V)) & \xrightarrow{-V_\ell(c_2)} & V_\ell(\text{CH}^2(V)) & \xrightarrow{V_\ell(\lambda^2)} & H^3(V, \mathbf{Q}_\ell(2)) \\ \uparrow f_* & & \xrightarrow{V_\ell(\det)} & & \uparrow f_* \\ V_\ell(\mathbf{K}_0(D)) & \xrightarrow{V_\ell(\det)} & V_\ell(\text{Pic}(D)) & \xrightarrow{V_\ell(\lambda^1)} & H^1(D, \mathbf{Q}_\ell(1)) \end{array}$$

commutes, where the right-hand side vertical arrow is the map (3.14) and the left-hand side vertical arrow is induced by the composition of the canonical map $\mathbf{K}_0(D) \rightarrow \mathbf{G}_0(D)$, which sends the rank 0 subgroup of $\mathbf{K}_0(D)$ to $F_{d-1} \mathbf{G}_0(D)$ (see [38, X, Corollaire 1.3.3]), with $f_ : F_{d-1} \mathbf{G}_0(D) \rightarrow F_{d-1} \mathbf{G}_0(V)$.*

We stress that D is not assumed to be reduced in Lemma 3.6.

Proof. – Let $\widetilde{\mathbf{K}}_0(D) = \text{Ker}(\text{rk} : \mathbf{K}_0(D) \rightarrow \mathbf{Z})$ and $\text{Gr}_\gamma^1 \mathbf{K}_0(D) = \widetilde{\mathbf{K}}_0(D)/\text{SK}_0(D)$ (a piece of notation justified by the fact that $\text{SK}_0(D)$ and $\widetilde{\mathbf{K}}_0(D)$ form the beginning of the γ -filtration on $\mathbf{K}_0(D)$). As the canonical map $\mathbf{K}_0(D) \rightarrow \mathbf{G}_0(D)$ sends $\widetilde{\mathbf{K}}_0(D)$ to $F_{d-1} \mathbf{G}_0(D)$ and $\text{SK}_0(D)$ to $F_{d-2} \mathbf{G}_0(D)$ (see [38, X, Corollaire 1.3.3]), there is an induced

map $f_* : V_\ell(\mathrm{Gr}_Y^1 \mathbf{K}_0(D)) \rightarrow V_\ell(\mathrm{Gr}_F^2 \mathbf{K}_0(V))$ and it suffices to prove the commutativity of the diagram

$$(3.16) \quad \begin{array}{ccccc} V_\ell(\mathrm{Gr}_F^2 \mathbf{K}_0(V)) & \xrightarrow[\sim]{-V_\ell(c_2)} & V_\ell(\mathrm{CH}^2(V)) & \xrightarrow{V_\ell(\lambda^2)} & H^3(V, \mathbf{Q}_\ell(2)) \\ \uparrow f_* & & \uparrow \text{dotted} & & \uparrow f_* \\ V_\ell(\mathrm{Gr}_Y^1 \mathbf{K}_0(D)) & \xrightarrow[\sim]{V_\ell(\det)} & V_\ell(\mathrm{Pic}(D)) & \xrightarrow{V_\ell(\lambda^1)} & H^1(D, \mathbf{Q}_\ell(1)), \end{array}$$

without the dotted arrow. We note that the leftmost horizontal arrows of (3.16) are isomorphisms; their inverses are induced by the map $\varphi^2 : \mathrm{CH}^2(V) \rightarrow \mathrm{Gr}_F^2 \mathbf{K}_0(V)$ of (2.4) and by the map $\mathrm{Pic}(D) \rightarrow \mathrm{Gr}_Y^1 \mathbf{K}_0(D)$ sending the class of a Cartier divisor Z on D to the class of $[\mathcal{O}_D(Z)] - [\mathcal{O}_D] \in \widetilde{\mathbf{K}}_0(D)$.

Let us complete this diagram with a dotted arrow induced by the composition of the canonical map $\mathrm{Pic}(D) \rightarrow \mathrm{CH}_{d-1}(D)$ (see [31, §2.1]) with the push-forward $f_* : \mathrm{CH}_{d-1}(D) \rightarrow \mathrm{CH}_{d-1}(V)$.

When D is smooth, the right half of (3.16) commutes by [10, Proposition 3.3, Proposition 3.6] and the left half by the description of the inverses of the horizontal arrows. Thus (3.16) commutes in this case.

In general, let us choose a family $(D_j)_{j \in J}$ of smooth projective varieties of pure dimension d , and for each $j \in J$, a morphism $v_j : D_j \rightarrow D$ and an element $n_j \in \mathbf{Z}_\ell$, such that the equality of d -cycles with coefficients in \mathbf{Z}_ℓ

$$(3.17) \quad [D] = \sum_{j \in J} n_j v_{j*} [D_j]$$

holds. When $\dim(D) \leq 2$ (which will be the case when we apply the lemma), one can choose the D_j to be desingularisations of the irreducible components of D^{red} and the n_j to be the multiplicities, in D , of these irreducible components. In arbitrary dimension, such D_j, v_j and n_j exist by the Gabber–de Jong alteration theorem [48, Theorem 2.1] applied to the irreducible components of D^{red} .

For $j \in J$, let $f_j = f \circ v_j : D_j \rightarrow V$. As D_j is smooth, we have already seen that the outer square of (3.16) with D and f replaced with D_j and f_j commutes. In order to show that the outer square of (3.16) itself commutes, it therefore suffices, by the contravariant functoriality of the lower row of (3.16), to check the equality $f_* = \sum_{j \in J} n_j f_{j*} \circ v_j^*$ of maps $V_\ell(\mathrm{Gr}_Y^1 \mathbf{K}_0(D)) \rightarrow V_\ell(\mathrm{Gr}_F^2 \mathbf{K}_0(V))$ and the same equality of maps $H^1(D, \mathbf{Q}_\ell(1)) \rightarrow H^3(V, \mathbf{Q}_\ell(2))$. Let us set $\mathrm{Gr}_i^F \mathbf{G}_0(D) = F_i \mathbf{G}_0(D) / F_{i-1} \mathbf{G}_0(D)$ and denote by $\kappa_D : \mathrm{Gr}_Y^1 \mathbf{K}_0(D) \rightarrow \mathrm{Gr}_{d-1}^F \mathbf{G}_0(D)$ the map induced by the canonical map $\mathbf{K}_0(D) \rightarrow \mathbf{G}_0(D)$. Coming back to the definition of f_* in the two contexts, we now see that it is enough to check the equalities

$$(3.18) \quad \kappa_D = \sum_{j \in J} n_j v_{j*} \circ \kappa_{D_j} \circ v_j^* : V_\ell(\mathrm{Gr}_Y^1 \mathbf{K}_0(D)) \rightarrow V_\ell(\mathrm{Gr}_{d-1}^F \mathbf{G}_0(D))$$

and

$$(3.19) \quad \kappa_D^{\acute{e}t} = \sum_{j \in J} n_j v_{j*} \circ \kappa_{D_j}^{\acute{e}t} \circ v_j^* : H^1(D, \mathbf{Q}_\ell(1)) \rightarrow H_{2d-1}(D, \mathbf{Q}_\ell(1-d)).$$

The $\mathbf{K}_0(D)$ -module structure of $\mathbf{G}_0(D)$ induces for any i a cap product operation $\mathrm{Gr}_Y^1 \mathbf{K}_0(D) \times \mathrm{Gr}_i^F \mathbf{G}_0(D) \rightarrow \mathrm{Gr}_{i-1}^F \mathbf{G}_0(D)$ (see [38, X, Corollaire 1.3.3]). Letting $[\mathcal{O}_D]$

denote the class of \mathcal{O}_D in $\mathrm{Gr}_d^F \mathbf{G}_0(D)$, we have $\kappa_D(x) = x \cap [\mathcal{O}_D]$ for any $x \in \mathrm{Gr}_v^1 \mathbf{K}_0(D)$; moreover (3.17) implies the equality $[\mathcal{O}_D] = \sum_{j \in J} n_j v_{j*} [\mathcal{O}_{D_j}]$ in $\mathrm{Gr}_d^F \mathbf{G}_0(D) \otimes_{\mathbf{Z}} \mathbf{Z}_\ell$ (see [31, Example 15.1.5]). In view of the projection formula [38, IV, (2.11.1.2)], we deduce (3.18). Similarly, the definition of $\kappa_D^{\acute{e}t}$, the equality obtained by applying cl to (3.17) and the projection formula [63, Proposition 4.2] together imply (3.19). \square

Let us finally start the proof of Proposition 3.5. As X is smooth over k , the morphism h gives rise to a push-forward map $h_* : H^3(X'_k, \mathbf{Q}_\ell(2)) \rightarrow H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2))$ (see (3.14)), satisfying $h_* \circ h^* = \mathrm{Id}$ on $H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2))$, since $h_*(h^*x \cap \mathrm{cl}([X'_k])) = x \cap \mathrm{cl}(h_*[X'_k]) = x \cap \mathrm{cl}([X_{\bar{k}}])$ for $x \in H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2))$ (see [63, Proposition 4.2]).

Let $K = \mathrm{Ker}(h_* : H^3(X'_k, \mathbf{Q}_\ell(2)) \rightarrow H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2)))$. We obtain a decomposition

$$(3.20) \quad H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2)) \oplus K \xrightarrow{\sim} H^3(X'_k, \mathbf{Q}_\ell(2)).$$

A second decomposition of the right-hand side can be obtained using the formula for the étale cohomology of the blow-up of a regularly immersed closed subscheme [75, Proposition 2.2.2.1] applied to the blow-ups appearing in diagram (3.3). This yields a canonical isomorphism

$$(3.21) \quad \bigoplus_{c_j=2} H^1((Z_j)_{\bar{k}}, \mathbf{Q}_\ell(1)) \xrightarrow{\sim} H^3(X'_k, \mathbf{Q}_\ell(2))$$

even though both X'_k and $(Z_j)_{\bar{k}}$ may fail to be regular.

Let Z'_j denote the normalization of $(Z_j)_{\bar{k}}^{\mathrm{red}}$ and $v_j : Z'_j \rightarrow (Z_j)_{\bar{k}}$ the natural morphism. The normality of Z_j implies that $(Z_j)_{\bar{k}}$ is geometrically unibranch and hence that v_j is universally bijective (see [35, Proposition 6.15.6, Proposition 6.15.5]). We deduce that $v_j^* : H^1((Z_j)_{\bar{k}}, \mathbf{Q}_\ell(1)) \rightarrow H^1(Z'_j, \mathbf{Q}_\ell(1))$ is an isomorphism (see [39, VIII, Corollaire 1.2]).

Letting $L = V_\ell(A(G_{\bar{k}}^0))$, we now consider the diagram of isomorphisms

$$(3.22) \quad \begin{array}{ccc} H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2)) \oplus K & \xrightarrow{\sim} & H^3(X'_k, \mathbf{Q}_\ell(2)) \leftarrow \sim \bigoplus_{c_j=2} H^1((Z_j)_{\bar{k}}, \mathbf{Q}_\ell(1)) \\ & & \downarrow v_j^* \wr \\ V_\ell((\mathbf{CH}_{X/k}^2)_{\bar{k}}^0) \oplus L & & \bigoplus_{c_j=2} H^1(Z'_j, \mathbf{Q}_\ell(1)) \\ \downarrow \wr & & \downarrow \wr \\ V_\ell(A((\mathbf{CH}_{X/k}^2)_{\bar{k}}^0)) \oplus L & \xrightarrow{\sim} & V_\ell(A((\mathbf{CH}_{X'/k}^2)_{\bar{k}}^0)) \leftarrow \sim \bigoplus_{c_j=2} V_\ell(A((\mathbf{Pic}_{Z_j/k}^0)_{\bar{k}})), \end{array}$$

whose upper horizontal arrows are (3.20) and (3.21), whose lower horizontal arrows stem from (3.12), whose left vertical isomorphism results from Theorem 3.1 (ii) and whose lower right vertical isomorphism is the composition of the canonical isomorphism $V_\ell(\mathrm{Pic}(Z'_j)) \xrightarrow{\sim} V_\ell(A((\mathbf{Pic}_{Z_j/k}^0)_{\bar{k}}))$ coming from §1.4 and (1.4) with the Kummer isomorphism $H^1(Z'_j, \mathbf{Q}_\ell(1)) \xrightarrow{\sim} V_\ell(\mathrm{Pic}(Z'_j))$.

Let m_j denote the multiplicity of $(Z_j)_{k_p}$, i.e., the length of its generic local ring, and θ_j the canonical principal polarization of $\mathbf{Pic}_{Z_j/k}^0$ (see §1.4.2).

- LEMMA 3.7. – 1. Diagram (3.22) transports the opposite of the cup product pairing on $H^3(X'_k, \mathbf{Q}_\ell(2))$ to the pairing on $\bigoplus_{c_j=2} V_\ell(A((\mathbf{Pic}^0_{Z_j/k})_{\bar{k}}))$ defined as the orthogonal sum of the $\mathbf{Q}_\ell(1)$ -valued Weil pairings associated with the polarizations $m_j\theta_j$.
2. Diagram (3.22) transports $H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2))$ to $V_\ell((\mathbf{CH}^2_{X/k})^0_{\bar{k}})$ and K to L .
3. The isomorphism $H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2)) = V_\ell((\mathbf{CH}^2_{X/k})^0_{\bar{k}})$ resulting from (ii) coincides with the one induced by Bloch’s ℓ -adic Abel-Jacobi map and by the identification between $V_\ell((\mathbf{CH}^2_{X/k})^0_{\bar{k}})$ and $V_\ell(\mathbf{CH}^2(X_{\bar{k}}))$ that stems from Theorem 3.1 (iv), (v).

Proof. – Let us consider the commutative diagram

$$\begin{array}{ccccc}
 \bigoplus_{c_j=2} H^1((Z_j)_{\bar{k}}, \mathbf{Q}_\ell(1)) & \xrightarrow{v_j^*} & \bigoplus_{c_j=2} H^1(Z'_j, \mathbf{Q}_\ell(1)) & & \\
 \downarrow \wr & & \downarrow \wr & & \\
 \bigoplus_{c_j=2} V_\ell(\mathbf{Pic}((Z_j)_{\bar{k}})) & \xrightarrow{v_j^*} & \bigoplus_{c_j=2} V_\ell(\mathbf{Pic}(Z'_j)) & & \\
 \parallel & & \parallel & & \\
 \bigoplus_{c_j=2} V_\ell(\mathbf{Pic}_{Z_j/k}) & \xleftarrow{\sim} & \bigoplus_{c_j=2} V_\ell(\mathbf{Pic}^0_{Z_j/k}) & \longrightarrow & \bigoplus_{c_j=2} V_\ell(A((\mathbf{Pic}^0_{Z_j/k})_{\bar{k}})) \\
 \parallel & & \parallel & & \parallel \\
 V_\ell(\mathbf{CH}^2_{X'/k}) & \xleftarrow{\sim} & V_\ell((\mathbf{CH}^2_{X'/k})^0) & \longrightarrow & V_\ell(A((\mathbf{CH}^2_{X'/k})^0_{\bar{k}})),
 \end{array}
 \tag{3.23}$$

in which the unlabeled horizontal arrows are the obvious ones (the bottom leftward arrow being an isomorphism in view of (3.12)), the top vertical arrows are the Kummer isomorphisms, the middle vertical isomorphisms come from §1.4 and (1.4), and the bottom vertical isomorphisms are induced by (3.12).

Since the top horizontal arrow of this diagram is an isomorphism, all of the maps appearing in (3.23) have to be isomorphisms.

We note that as a consequence of the projection formula [63, Proposition 4.2] and of the equality of cycles $[(Z_j)_{\bar{k}}] = m_j v_{j*}[Z'_j]$, the top horizontal isomorphism of (3.23) transports the cup product pairing on $H^1((Z_j)_{\bar{k}}, \mathbf{Q}_\ell(1))$ to the cup product pairing on $H^1(Z'_j, \mathbf{Q}_\ell(1))$ multiplied by m_j .

As on the other hand (3.23) transports the Weil pairing on $V_\ell(A((\mathbf{Pic}^0_{Z_j/k})_{\bar{k}})) = V_\ell(\mathbf{Pic}^0_{Z'_j/\bar{k}})$ (see (1.4)) to the cup product pairing on $H^1(Z'_j, \mathbf{Q}_\ell(1))$, we see that Lemma 3.7 (i) amounts to the assertion that (3.21) transports the orthogonal sum of the cup product pairings on $H^1((Z_j)_{\bar{k}}, \mathbf{Q}_\ell(1))$ to the opposite of the cup product pairing on $H^3(X'_k, \mathbf{Q}_\ell(2))$. When the Z_j are smooth, this is shown in [7, (2.9)]; the same argument applies in our setting.

Thus, it only remains to prove Lemma 3.7 (ii) and (iii). For this, it suffices to check the commutativity of the squares

$$\begin{array}{ccc}
 V_\ell(A((\mathbf{CH}^2_{X'/k})^0_{\bar{k}})) & \xrightarrow{\gamma'} & H^3(X'_k, \mathbf{Q}_\ell(2)) \\
 h^* \uparrow & & \uparrow h^* \\
 V_\ell(A((\mathbf{CH}^2_{X/k})^0_{\bar{k}})) & \xrightarrow{\gamma} & H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2))
 \end{array}
 \tag{3.24}$$

and

$$(3.25) \quad \begin{array}{ccc} V_\ell(A((\mathbf{CH}_{X'/k}^2)_{\bar{k}}^0)) & \xrightarrow{\sim} \xrightarrow{\gamma'} & H^3(X'_{\bar{k}}, \mathbf{Q}_\ell(2)) \\ h_* \downarrow & & \downarrow h_* \\ V_\ell(A((\mathbf{CH}_{X/k}^2)_{\bar{k}}^0)) & \xrightarrow{\sim} \xrightarrow{\gamma} & H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2)), \end{array}$$

where γ is the isomorphism constructed from Bloch's ℓ -adic Abel-Jacobi map for the smooth variety X (see the statement of Lemma 3.7 (iii)) and γ' is the isomorphism extracted from (3.21) and (3.23), and where the vertical arrows are those appearing in the upper and lower rows of (3.22).

The square (3.25) fits into a larger diagram

$$(3.26) \quad \begin{array}{ccccccc} \bigoplus_{c_j=2} V_\ell(\mathbf{K}_0((Z_j)_{\bar{k}})) & \xrightarrow{\sim} \xrightarrow{\det} & \bigoplus_{c_j=2} V_\ell(\text{Pic}((Z_j)_{\bar{k}})) & = & \bigoplus_{c_j=2} H^1((Z_j)_{\bar{k}}, \mathbf{Q}_\ell(1)) \\ \downarrow \alpha & & \parallel & & \parallel \\ V_\ell(\text{SK}_0(X'_{\bar{k}})) & \longrightarrow & V_\ell(\mathbf{CH}_{X'/k}^2) \xrightarrow{\beta'} & V_\ell(A((\mathbf{CH}_{X'/k}^2)_{\bar{k}}^0)) & \xrightarrow{\sim} \xrightarrow{\gamma'} & H^3(X'_{\bar{k}}, \mathbf{Q}_\ell(2)) \\ \downarrow h_* & & \downarrow h_* & \downarrow h_* & & \downarrow h_* \\ V_\ell(\text{SK}_0(X_{\bar{k}})) & \longrightarrow & V_\ell(\mathbf{CH}_{X/k}^2) \xrightarrow{\beta} & V_\ell(A((\mathbf{CH}_{X/k}^2)_{\bar{k}}^0)) & \xrightarrow{\sim} \xrightarrow{\gamma} & H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2)), \end{array}$$

in which the map α is induced by (3.6) (see also (3.7)), the map β' comes from the bottom row of (3.23), the map β is constructed in the same way as β' (legitimate thanks to Theorem 3.1 (v)) and the isomorphisms of the square of the top right corner all come from (3.21) and (3.23).

In order for the square in the bottom right corner to commute, it suffices that the outer square of the diagram commutes, since the other inner squares clearly commute. That is, fixing j such that $c_j = 2$ and letting α_j and $\alpha_j^{\acute{e}t}$ respectively denote the j -th component of α and of (3.21), we need only prove that the square

$$(3.27) \quad \begin{array}{ccc} V_\ell(\mathbf{K}_0((Z_j)_{\bar{k}})) & \xrightarrow{\sim} & H^1((Z_j)_{\bar{k}}, \mathbf{Q}_\ell(1)) \\ \downarrow h_* \circ \alpha_j & & \downarrow h_* \circ \alpha_j^{\acute{e}t} \\ V_\ell(\text{SK}_0(X_{\bar{k}})) & \longrightarrow & H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2)), \end{array}$$

whose horizontal arrows are extracted from (3.26), commutes.

Set $D_j = Z_j \times_{Y_j} X'$ and $E_j = Z_j \times_{Y_j} Y_{j+1}$, where the Y_j are the varieties appearing in diagram (3.3). Let $q_j : D_j \rightarrow Z_j$ and $p'_j : E_j \rightarrow Z_j$ denote the projections. Let $\iota_j : Z_j \hookrightarrow Y_j$, $\iota'_j : E_j \hookrightarrow Y_{j+1}$ and $\delta_j : D_j \hookrightarrow X'$ be the inclusions, so that $\iota_j, \iota'_j, \delta_j$ are regular closed immersions of codimensions $c_j, 1, 1$, respectively. Recall that $\alpha_j^{\acute{e}t} = (p_{j+1} \circ \dots \circ p_N)^* \circ \iota'_{j*} \circ p_j'^*$, where ι'_{j*} denotes the map given by cup product with the class of the Cartier divisor $(E_j)_{\bar{k}}$ in $H^2_{(E_j)_{\bar{k}}}((Y_{j+1})_{\bar{k}}, \mathbf{Q}_\ell(1))$ (see [75, §2.1]) composed with the map

$$H^3_{(E_j)_{\bar{k}}}((Y_{j+1})_{\bar{k}}, \mathbf{Q}_\ell(2)) \rightarrow H^3((Y_{j+1})_{\bar{k}}, \mathbf{Q}_\ell(2))$$

forgetting the support. As E_j pulls back, as a Cartier divisor, to D_j , we deduce that

$$(3.28) \quad \alpha_j^{\acute{e}t} = \delta_{j*} \circ q_j^*,$$

where δ_{j*} is again defined as in *loc. cit.*, §2.1.

Similarly, recall that α_j is given by $x \mapsto (p_{j+1} \circ \dots \circ p_N)^*(l'_{j*} p_j'^* x \otimes [\mathcal{O}_{Y_{j+1}}(E_j)]) = ((p_{j+1} \circ \dots \circ p_N)^* l'_{j*} p_j'^* x) \otimes [\mathcal{O}_{X'}(D_j)]$ (see (2.3)). Noting that the morphisms $p_{j+1} \circ \dots \circ p_N$ and l'_j are Tor-independent (indeed one has $\text{Tor}_i^A(A/fA, B) = 0$ for any $i > 0$, any commutative ring A , any A -algebra B and any $f \in A$ such that neither f nor its image in B is a zero divisor), the base change theorem [64, Theorem 3.10.3] allows us to rewrite this as

$$(3.29) \quad \alpha_j(x) = (\delta_{j*} \circ q_j^*)(x) \otimes [\mathcal{O}_{X'}(D_j)]$$

for any $x \in V_\ell(\mathbf{K}_0((Z_j)_{\bar{k}}))$.

Let $\alpha'_j : V_\ell(\mathbf{K}_0((Z_j)_{\bar{k}})) \rightarrow V_\ell(\mathbf{SK}_0(X'_k))$ be given by $\alpha'_j(x) = (\delta_{j*} \circ q_j^*)(x)$. In view of (3.28) and of the contravariant functoriality of the first row of (3.26), we deduce from Lemma 3.6 applied to $h \circ \delta_j : D_j \rightarrow X$ that the square obtained by replacing, in (3.27), the left-hand side vertical arrow $h_* \circ \alpha_j$ with $h_* \circ \alpha'_j$ commutes. On the other hand, it follows from (3.29) that the map $h_* \circ \alpha_j - h_* \circ \alpha'_j$ takes its values in $V_\ell(F^3 \mathbf{K}_0(X_{\bar{k}}))$, since $\text{rk}([\mathcal{O}_{X'}(D_j)] - [\mathcal{O}_{X'}]) = 0$ (see [38, X, Corollaire 1.3.3]). Now the lower horizontal map of (3.27) vanishes on $V_\ell(F^3 \mathbf{K}_0(X_{\bar{k}}))$ since it factors through c_2 ; we conclude that the square (3.27) itself commutes, and therefore so does (3.25).

Let us turn to (3.24). We introduce a desingularisation $\pi : X'' \rightarrow X'_k$ of X'_k (which exists by Cossart and Piltant [23]) and consider the square

$$(3.30) \quad \begin{array}{ccc} V_\ell(A((\mathbf{CH}^2_{X'/k})^0_{\bar{k}})) & \xrightarrow{\sim} \gamma' & H^3(X'_k, \mathbf{Q}_\ell(2)) \\ \pi^* \downarrow & & \downarrow \pi^* \\ V_\ell(A((\mathbf{CH}^2_{X''/\bar{k}})^0)) & \xrightarrow{\sim} \gamma'' & H^3(X'', \mathbf{Q}_\ell(2)), \end{array}$$

where γ'' is constructed from Bloch’s ℓ -adic Abel-Jacobi map for the smooth variety X'' in the same way as γ for X . One verifies the commutativity of the square (3.30) by proceeding exactly as we did with (3.25), that is, by reducing to Lemma 3.6 using the diagram obtained by replacing, in (3.26), all occurrences of X with X'' and all occurrences of h_* with π^* , and using the equalities obtained by replacing, in (3.28) and (3.29) and in their proofs, α_j and $\alpha_j^{\text{ét}}$ with $\pi^* \circ \alpha_j$ and $\pi^* \circ \alpha_j^{\text{ét}}$, and D_j, q_j, δ_j with D''_j, q''_j, δ''_j , where $D''_j = Z_j \times_{Y_j} X''$ and where $q''_j : D''_j \rightarrow Z_j$ and $\delta''_j : D''_j \hookrightarrow X''$ denote the projections.

All the \bar{k} -varieties in sight and all the \bar{k} -morphisms between them are defined over a common subfield of \bar{k} that is finitely generated over the prime field. Their ℓ -adic cohomology groups are thus endowed with a weight filtration (see [51, §2]). As the Z'_j are smooth and projective, the groups $H^1(Z'_j, \mathbf{Q}_\ell(1))$ are pure of weight -1 . In view of the isomorphisms (3.22), we deduce that the group $H^3(X'_k, \mathbf{Q}_\ell(2))$ is pure of weight -1 as well. On the other hand, the kernel of the right-hand side vertical map of (3.30) has weights < -1 , as follows from cohomological descent and Deligne’s theorem on the Weil conjectures (*op. cit.*, §9; proper smooth hypercoverings of X'_k that start with π exist by [52, Theorem 4.1]). Hence the right-hand side vertical map of (3.30) is injective.

This injectivity, the commutativity of (3.30) and the commutativity of the square

$$(3.31) \quad \begin{array}{ccc} V_\ell(A((\mathbf{CH}_{X''/\bar{k}}^2)^0)) & \xrightarrow[\sim]{\gamma''} & H^3(X'', \mathbf{Q}_\ell(2)) \\ \pi^* \circ h^* \uparrow & & \uparrow \pi^* \circ h^* \\ V_\ell(A((\mathbf{CH}_{X/k}^2)^0)) & \xrightarrow[\sim]{\gamma} & H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2)) \end{array}$$

(which holds by functoriality of Bloch's Abel-Jacobi map [10, Proposition 3.5]) together imply that (3.24) commutes. This concludes the proof of Lemma 3.7. \square

We resume the proof of Proposition 3.5. Consider the diagram

$$(3.32) \quad (\mathbf{CH}_{X/k}^2)^0_{\bar{k}} \times A(G_{\bar{k}}^0) \xrightarrow{\sim} A((\mathbf{CH}_{X'/k}^2)^0_{\bar{k}}) \xleftarrow{\sim} \prod_{c_j=2} A((\mathbf{Pic}_{Z_j/k}^0)_{\bar{k}})$$

of isomorphisms of abelian varieties stemming from (3.12) and whose ℓ -adic Tate modules appear on the bottom line of (3.22). The product of the polarizations $m_j \theta_j$ on the right-hand side of (3.32) induces a polarization λ on the left-hand side $(\mathbf{CH}_{X/k}^2)^0_{\bar{k}} \times A(G_{\bar{k}}^0)$ of (3.32). Let us view the Weil pairing of λ as a $\mathbf{Q}_\ell(1)$ -valued pairing on $H^3(X'_{\bar{k}}, \mathbf{Q}_\ell(2))$ thanks to (3.22). By Lemma 3.7 (i), it is equal to the opposite of the cup product pairing on $H^3(X'_{\bar{k}}, \mathbf{Q}_\ell(2))$. Since, by the projection formula [63, Proposition 4.2], the decomposition (3.20) is orthogonal with respect to the cup product, it follows from Lemma 3.7 (ii) that λ is a product polarization on $(\mathbf{CH}_{X/k}^2)^0_{\bar{k}} \times A(G_{\bar{k}}^0)$. Since the restriction of the cup product pairing on $H^3(X'_{\bar{k}}, \mathbf{Q}_\ell(2))$ to $H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2))$ coincides with the cup product pairing on $H^3(X_{\bar{k}}, \mathbf{Q}_\ell(2))$, it follows from Lemma 3.7 (iii) that the restriction of λ to $(\mathbf{CH}_{X/k}^2)^0_{\bar{k}}$ is the canonical principal polarization defined in Theorem 3.1 (vi).

By a theorem of Debarre [24, Corollary 2], polarized abelian varieties can be written in a unique way as a product of indecomposable polarized abelian varieties. As the $(A((\mathbf{Pic}_{Z_j/k}^0)_{\bar{k}}), m_j \theta_j)$ are indecomposable or trivial (since so are the $(A((\mathbf{Pic}_{Z_j/k}^0)_{\bar{k}}), \theta_j)$), we deduce the existence of a partition $\{j \mid c_j = 2\} = J \sqcup J'$ such that (3.32) induces isomorphisms $\prod_{j \in J} A((\mathbf{Pic}_{Z_j/k}^0)_{\bar{k}}) \xrightarrow{\sim} (\mathbf{CH}_{X/k}^2)^0_{\bar{k}}$ and $\prod_{j \in J'} A((\mathbf{Pic}_{Z_j/k}^0)_{\bar{k}}) \xrightarrow{\sim} A(G_{\bar{k}}^0)$. Since λ restricts to a principal polarization on $(\mathbf{CH}_{X/k}^2)^0_{\bar{k}}$, we see that $m_j = 1$ for all the $j \in J$ such that $A((\mathbf{Pic}_{Z_j/k}^0)_{\bar{k}})$ is non-zero.

Thus, the product of the polarizations θ_j on the right-hand side of (3.32) induces on $(\mathbf{CH}_{X/k}^2)^0_{\bar{k}} \times A(G_{\bar{k}}^0)$ a polarization which is at the same time a principal polarization and the product of two polarizations, and which is therefore the product of two principal polarizations; moreover, the first of these coincides with the canonical principal polarization of Theorem 3.1 (vi). Proposition 3.5 is proved. \square

Now that Proposition 3.5 is proved, we let J_1 (resp. J_2 , resp. J_3) be the set of indices j such that $c_j = 2$ and the curve Z_j is smooth over k (resp. such that $c_j = 2$ and Z_j is not smooth over k , resp. such that $c_j = 3$), and we proceed to show that the $(Z_j)_{j \in J_2}$ do not contribute to $(\mathbf{CH}_{X/k}^2)^0$.

LEMMA 3.8. – *The map $\prod_{j \in J_2} \mathbf{Pic}_{Z_j/k}^0 \rightarrow (\mathbf{CH}_{X/k}^2)^0$ induced by (3.12) vanishes.*

Proof. – We fix $j \in J_2$. Let $a : \mathbf{Pic}_{Z_j/k}^0 \rightarrow (\mathbf{CH}_{X/k}^2)^0$ and $b : \mathbf{Pic}_{Z_j/k}^0 \rightarrow G^0$ denote the maps induced by (3.12). Let us assume that $a \neq 0$ and derive a contradiction. When $a \neq 0$, we claim that $\mathbf{Pic}_{Z_j/k}^0 = \text{Ker}(a) \times \text{Ker}(b)$, that $\text{Ker}(a)$ is affine and that $\text{Ker}(b)$ is a non-trivial abelian variety; Corollary 1.8 then provides the desired contradiction. It thus suffices to prove the claim. To this end, it is enough to check that

$$(\mathbf{Pic}_{Z_j/k}^0)_{k_p} = \text{Ker}(a_{k_p}) \times \text{Ker}(b_{k_p}),$$

that $\text{Ker}(a_{k_p})$ is affine and that $\text{Ker}(b_{k_p})$ is a non-trivial abelian variety, as these three properties descend to k .

By functoriality, the map b induces maps $A(b_{k_p}) : A((\mathbf{Pic}_{Z_j/k}^0)_{k_p}) \rightarrow A(G_{k_p}^0)$ and $L(b_{k_p}) : L((\mathbf{Pic}_{Z_j/k}^0)_{k_p}) \rightarrow L(G_{k_p}^0)$. As $(\mathbf{CH}_{X/k}^2)_{k_p}^0$ is an abelian variety (see Theorem 3.1 (ii)), we have $L(a_{k_p}) = 0$ and the map $A(a_{k_p})$ can be viewed as a map

$$A(a_{k_p}) : A((\mathbf{Pic}_{Z_j/k}^0)_{k_p}) \rightarrow (\mathbf{CH}_{X/k}^2)_{k_p}^0$$

through which a_{k_p} factors, so that our assumption that $a_{k_p} \neq 0$ implies that $A(a_{k_p}) \neq 0$. On the other hand, the map $L(b_{k_p})$ is a closed immersion since $a \times b$ is one and $L(a_{k_p}) = 0$.

Proposition 3.5 allows us to view $A((\mathbf{Pic}_{Z_j/k}^0)_{k_p})$ as a principally polarized direct factor of the product of principally polarized abelian varieties $(\mathbf{CH}_{X/k}^2)_{k_p}^0 \times A(G_{k_p}^0)$ over k_p , through $A(a_{k_p}) \times A(b_{k_p})$. As the decomposition of a principally polarized abelian variety into its indecomposable factors is unique, as $A((\mathbf{Pic}_{Z_j/k}^0)_{k_p})$ is itself indecomposable (see §1.4.2), and as $A(a_{k_p}) \neq 0$, necessarily $A(b_{k_p}) = 0$ and $A(a_{k_p})$ is a closed immersion, so that $\text{Ker}(a_{k_p}) = L((\mathbf{Pic}_{Z_j/k}^0)_{k_p})$ (see (1.1)).

All in all, the exact sequences (1.1) fit into a commutative diagram

$$(3.33) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker}(a_{k_p}) & \longrightarrow & (\mathbf{Pic}_{Z_j/k}^0)_{k_p} & \longrightarrow & A((\mathbf{Pic}_{Z_j/k}^0)_{k_p}) \longrightarrow 0 \\ & & \downarrow & & \downarrow b_{k_p} & & \downarrow 0 \\ 0 & \longrightarrow & L(G_{k_p}^0) & \longrightarrow & G_{k_p}^0 & \longrightarrow & A(G_{k_p}^0) \longrightarrow 0. \end{array}$$

An isomorphism $\text{Ker}(b_{k_p}) \xrightarrow{\sim} A((\mathbf{Pic}_{Z_j/k}^0)_{k_p})$ and then all of the desired statements now result from this diagram, in view of the remark that the snake homomorphism is trivial since it goes from an abelian variety to an affine group. \square

REMARK 3.9. – At the end of the proof of Proposition 3.5 (penultimate paragraph), we had seen, as a by-product of [24], that for any $j \in \{1, \dots, N\}$ such that $c_j = 2$, if the map $A((\mathbf{Pic}_{Z_j/k}^0)_{\bar{k}}) \rightarrow (\mathbf{CH}_{X/k}^2)_{\bar{k}}^0$ induced by (3.12) does not vanish, then the curve Z_j is geometrically reduced. Lemma 3.8 reproves this fact by other means, in the course of showing that Z_j is even smooth. Instead of reproving it, it would have been possible to use the geometric reducedness of Z_j to simplify the proof of Lemma 3.8, inasmuch as Corollary 1.8 could have been applied over k_p .

Proof of Theorem 3.1 (vii). – In view of Lemma 3.8, we may consider the quotient H of G by its subgroup scheme $\prod_{j \in J_2} \mathbf{Pic}_{Z_j/k}^0$.

Thanks to the exact sequences $0 \rightarrow \mathbf{Pic}_{Z_j/k}^0 \rightarrow \mathbf{Pic}_{Z_j/k} \rightarrow \mathbf{Z}_{Z_j/k} \rightarrow 0$ given by Proposition 1.6 for $j \in J_2$, we deduce from (3.12) an isomorphism

$$(3.34) \quad \mathbf{CH}_{X/k}^2 \times H \xleftarrow{\sim} \mathbf{Z} \times \prod_{j \in J_1} \mathbf{Pic}_{Z_j/k} \times \prod_{j \in J_2 \cup J_3} \mathbf{Z}_{Z_j/k}.$$

Let B be the disjoint union of \mathbf{P}_k^1 , of the curves $\mathbf{P}_{\pi_0(Z_j/k)}^1$ for all $j \in J_2 \cup J_3$, and of the curves Z_j for all $j \in J_1$. It is a smooth projective curve over k , and it has the property that

$$\mathbf{Pic}_{B/k} \simeq \mathbf{Z} \times \prod_{j \in J_1} \mathbf{Pic}_{Z_j/k} \times \prod_{j \in J_2 \cup J_3} \mathbf{Z}_{Z_j/k}$$

since $\mathbf{Pic}_{\mathbf{P}_{\pi_0(Z_j/k)}^1/k} \simeq \text{Res}_{\pi_0(Z_j/k)/k}(\mathbf{Z}) \simeq \mathbf{Z}_{Z_j/k}$ for $j \in J_2 \cup J_3$ by Proposition 1.1. The isomorphism $\mathbf{Pic}_{B/k} \xrightarrow{\sim} \mathbf{CH}_{X/k}^2 \times H$ deduced from (3.34) realizes $\mathbf{CH}_{X/k}^2$ as a principally polarized direct factor of $\mathbf{Pic}_{B/k}$ by Proposition 3.5, as desired. \square

The proof of Theorem 3.1 is now complete.

3.3. Blow-ups

The next proposition, which relies on arguments already used in the proof of Theorem 3.1, allows one to compute $\mathbf{CH}_{X/k}^2$ in concrete situations.

PROPOSITION 3.10. – *Let X be a smooth projective \bar{k} -rational threefold over k , let $i : Y \rightarrow X$ be the inclusion of a smooth closed subvariety of pure codimension c , and let $p : X' \rightarrow X$ be its blow-up. Then the formula (2.8) induces an isomorphism*

$$(3.35) \quad \begin{aligned} \mathbf{CH}_{X/k}^2 \times \mathbf{Pic}_{Y/k} &\xrightarrow{\sim} \mathbf{CH}_{X'/k}^2 && \text{if } c = 2 \\ (\text{resp. } \mathbf{CH}_{X/k}^2 \times \mathbf{Z}_{Y/k}) &\xrightarrow{\sim} \mathbf{CH}_{X'/k}^2 && \text{if } c = 3), \end{aligned}$$

respecting the principal polarizations furnished by Theorem 3.1 (applied to X and to X') and by §1.4.2 (applied to Y).

Proof. – The isomorphism (2.8), Corollary 1.5 and Remark 1.2 (ii) yield canonical isomorphisms $\mathbf{K}_{0,X/k} \times \mathbf{K}_{0,Y/k}^{c-1} \xrightarrow{\sim} \mathbf{K}_{0,X'/k}$, $\mathbf{Pic}_{X/k} \times \mathbf{Z}_{Y/k} \xrightarrow{\sim} \mathbf{Pic}_{X'/k}$ and $\mathbf{Z}_{X/k} \xrightarrow{\sim} \mathbf{Z}_{X'/k}$. Identifying $(\text{rk}, \det) : \mathbf{K}_{0,X'/k} \rightarrow \mathbf{Z}_{X'/k} \times \mathbf{Pic}_{X'/k}$ in terms of these decompositions and using Proposition 2.4, we obtain an isomorphism

$$(3.36) \quad \begin{aligned} \mathbf{SK}_{0,X/k,\text{fppf}} \times \mathbf{Pic}_{Y/k,\text{fppf}} &\xrightarrow{\sim} \mathbf{SK}_{0,X'/k,\text{fppf}} && \text{if } c = 2 \\ (\text{resp. } \mathbf{SK}_{0,X/k,\text{fppf}} \times \mathbf{Z}_{Y/k}) &\xrightarrow{\sim} \mathbf{SK}_{0,X'/k,\text{fppf}} && \text{if } c = 3). \end{aligned}$$

If l/k is a finite extension and $x \in (X \setminus Y)(l)$, one has $p^*[\mathcal{O}_x] = [\mathcal{O}_{p^{-1}(x)}] \in \mathbf{K}_0(X'_l)$. It follows that $p^*v_X(1) = v_{X'}(1) \in \mathbf{SK}_{0,X'/k,\text{fppf}}(k) \subseteq \mathbf{SK}_{0,X'/k,\text{fppf}}(l)$. We thus deduce from (3.36) the required isomorphism (3.35) of k -group schemes.

If $c = 2$, considering the commutative diagram

$$(3.37) \quad \begin{array}{ccc} \text{Ab}^2(X_{k_p}) \times \mathbf{Pic}_{Y_{k_p}/k_p}^0 & \xrightarrow{\sim} & \text{Ab}^2(X'_{k_p}) \\ \downarrow \wr & & \downarrow \wr \\ (\mathbf{CH}_{X_{k_p}/k_p}^2)^0 \times \mathbf{Pic}_{Y_{k_p}/k_p}^0 & \xrightarrow{\sim} & (\mathbf{CH}_{X'_{k_p}/k_p}^2)^0 \end{array}$$

whose vertical arrows stem from Theorem 3.1 (vi), whose lower horizontal arrow is the above constructed isomorphism and whose upper horizontal arrow is that of [7, Lemma 2.10] concludes the proof, as the latter arrow respects the principal polarizations by [7, Lemma 2.10]. If $c = 3$, one can argue in the same way, using a diagram similar to (3.37) in which $\mathbf{Pic}_{Y_{k_p}/k_p}^0$ does not appear. \square

3.4. Obstructions to k -rationality

The most general obstruction to the k -rationality of a smooth projective \bar{k} -rational threefold obtained in this article is Theorem 3.1 (vii). In this subsection, we spell out concrete consequences of this theorem.

We recall that a Γ_k -module M is a *permutation* Γ_k -module if it is free of finite rank as a \mathbf{Z} -module and admits a \mathbf{Z} -basis that is permuted by the action of Γ_k , and that it is *stably of permutation* if there exists a Γ_k -equivariant isomorphism $M \oplus N_1 \simeq N_2$ for some permutation Γ_k -modules N_1 and N_2 .

If X is a smooth projective \bar{k} -rational threefold, we associate with any class $\alpha \in \mathrm{NS}^2(X_{\bar{k}})^{\Gamma_k} = (\mathbf{CH}_{X/k}^2 / (\mathbf{CH}_{X/k}^2)^0)(k)$ (see Theorem 3.1 (v)) its inverse image $(\mathbf{CH}_{X/k}^2)^\alpha$ in $\mathbf{CH}_{X/k}^2$. It is an fppf torsor under $(\mathbf{CH}_{X/k}^2)^0$, hence an étale torsor under $(\mathbf{CH}_{X/k}^2)^0$ by [68, III, Corollary 4.7, Remark 4.8 (a)]. We let $[(\mathbf{CH}_{X/k}^2)^\alpha] \in H^1(k, (\mathbf{CH}_{X/k}^2)^0)$ be its Galois cohomology class.

THEOREM 3.11. – *Let X be a smooth projective k -rational threefold over k . Then:*

- (i) *The Γ_k -module $\mathrm{NS}^2(X_{\bar{k}})$ is a direct factor of a permutation Γ_k -module.*
- (ii) *There exists an isomorphism $(\mathbf{CH}_{X/k}^2)^0 \simeq \mathbf{Pic}_{C/k}^0$ of principally polarized abelian varieties over k for some smooth projective curve C over k .*
- (iii) *For all smooth projective geometrically connected curves D of genus ≥ 2 over k , all morphisms $\psi : (\mathbf{CH}_{X/k}^2)^0 \rightarrow \mathbf{Pic}_{D/k}^0$ identifying $\mathbf{Pic}_{D/k}^0$ with a principally polarized direct factor of $(\mathbf{CH}_{X/k}^2)^0$ and all $\alpha \in \mathrm{NS}^2(X_{\bar{k}})^{\Gamma_k}$, there exists $d \in \mathbf{Z}$ such that $\psi_*[(\mathbf{CH}_{X/k}^2)^\alpha] = [\mathbf{Pic}_{D/k}^d] \in H^1(k, \mathbf{Pic}_{D/k}^0)$.*
- (iv) *For all elliptic curves E over k and all morphisms $\psi : (\mathbf{CH}_{X/k}^2)^0 \rightarrow E$ identifying E with a principally polarized direct factor of $(\mathbf{CH}_{X/k}^2)^0$, there exists a class $\eta \in H^1(k, E)$ such that for all $\alpha \in \mathrm{NS}^2(X_{\bar{k}})^{\Gamma_k}$, there exists $d \in \mathbf{Z}$ with $\psi_*[(\mathbf{CH}_{X/k}^2)^\alpha] = d\eta \in H^1(k, E)$.*

Proof. – Theorem 3.1 (vii) shows the existence of a smooth projective curve B over k such that $\mathbf{CH}_{X/k}^2$ is a principally polarized direct factor of $\mathbf{Pic}_{B/k}$.

We denote by $q : \mathbf{Pic}_{B/k} \rightarrow \mathbf{CH}_{X/k}^2$ the projection onto this direct factor and by $r : \mathbf{CH}_{X/k}^2 \rightarrow \mathbf{Pic}_{B/k}$ the inclusion of this direct factor.

Passing to the groups of connected components shows that $\mathrm{NS}^2(X_{\bar{k}})$ is a direct factor of $\mathbf{Z}_{B/k}(\bar{k})$, which is a permutation Γ_k -module, thus proving (i).

Passing to the identity components shows that $(\mathbf{CH}_{X/k}^2)^0$ is a principally polarized direct factor of $\mathbf{Pic}_{B/k}^0$. In view of the uniqueness of the decomposition of a principally polarized abelian variety as a product of indecomposable ones, and in view of the description of the indecomposable factors of $\mathbf{Pic}_{B/k}^0$ (see [7, §2.1]), there exists a union C of connected

components of B such that $(\mathbf{CH}_{X/k}^2)^0 \simeq \mathbf{Pic}_{C/k}^0$ as principally polarized abelian varieties over k , thus proving (ii).

Let us now fix D , ψ and α as in (iii). The composition $p := \psi \circ q^0 : \mathbf{Pic}_{B/k}^0 \rightarrow \mathbf{Pic}_{D/k}^0$ realizes $\mathbf{Pic}_{D/k}^0$ as a principally polarized direct factor of $\mathbf{Pic}_{B/k}^0$. All indecomposable principally polarized direct factors of $\mathbf{Pic}_{B/k}^0$ are of the form $\mathbf{Pic}_{B'/k}^0$ for some connected component B' of B (as the connected components of B are in bijection with those of B_{k_p} , this statement reduces to the case of a perfect field, for which see [7, §2.1]). Let B' be the connected component of B corresponding to $\mathbf{Pic}_{D/k}^0$. Since D is geometrically connected of genus ≥ 2 , we see that $\mathbf{Pic}_{D/k}^0$, hence also $\mathbf{Pic}_{B'/k}^0$, is geometrically indecomposable of dimension ≥ 2 , and it follows that B' is geometrically connected of genus ≥ 2 . By the precise form of the Torelli theorem [77, Théorèmes 1 et 2], after possibly replacing q with $-q$ (and p with $-p$) if D is not hyperelliptic, we can identify D and B' in such a way that p is the pull-back by the inclusion $i : D \simeq B' \hookrightarrow B$. Since $q : \mathbf{Pic}_{B/k} \rightarrow \mathbf{CH}_{X/k}^2$ realizes $\mathbf{CH}_{X/k}^2$ as a direct factor of $\mathbf{Pic}_{B/k}$, one can find $\beta \in \mathrm{NS}^1(B_{\bar{k}})^{\Gamma_k}$ with $q(\beta) = \alpha \in \mathrm{NS}^2(X_{\bar{k}})^{\Gamma_k}$. Letting $d := i^*\beta \in \mathrm{NS}^1(D_{\bar{k}}) \simeq \mathbf{Z}$, we obtain $\psi_*[(\mathbf{CH}_{X/k}^2)^\alpha] = p_*[\mathbf{Pic}_{B'/k}^\beta] = [\mathbf{Pic}_{D/k}^d] \in H^1(k, \mathbf{Pic}_{D/k}^0)$, which proves (iii).

Fix E and ψ as in (iv). Arguing as above shows that $p := \psi \circ q^0 : \mathbf{Pic}_{B/k}^0 \rightarrow E$ identifies E with $\mathbf{Pic}_{B'/k}^0$ for some connected component $i : B' \hookrightarrow B$ of B which is geometrically connected of genus 1. The genus 1 curve B' has a natural structure of $\mathbf{Pic}_{B'/k}^0$ -torsor, and we set $\eta := [B'] \in H^1(k, \mathbf{Pic}_{B'/k}^0) = H^1(k, E)$ its class. Let $\alpha \in \mathrm{NS}^2(X_{\bar{k}})^{\Gamma_k}$. Setting $s := i^* \circ r : \mathbf{CH}_{X/k}^2 \rightarrow \mathbf{Pic}_{B'/k}$ and $d := s_*\alpha \in \mathrm{NS}^1(B'_{\bar{k}}) \simeq \mathbf{Z}$, we get $\psi_*[(\mathbf{CH}_{X/k}^2)^\alpha] = s_*[(\mathbf{CH}_{X/k}^2)^\alpha] = [\mathbf{Pic}_{B'/k}^d] = d\eta \in H^1(k, \mathbf{Pic}_{B'/k}^0) = H^1(k, E)$, which completes the proof of (iv). \square

REMARKS 3.12. – (i) If X is a smooth projective k -rational variety and if k has characteristic 0, one can apply the weak factorization theorem [2, Theorem 0.3.1] to show that the Γ_k -module $\mathrm{NS}^2(X_{\bar{k}})$ is stably of permutation. This statement is stronger than Theorem 3.11 (i). We do not know if it holds if k has characteristic $p > 0$ and X has dimension ≥ 3 . We do not know either whether Theorem 3.11 (i) holds for smooth projective k -rational varieties of dimension ≥ 4 .

(ii) The geometric Néron-Severi group $\mathrm{NS}^1(X_{\bar{k}})$ of a smooth projective k -rational variety X is stably of permutation (see [66, Theorem 2.2] if X is a surface and [16, Proposition 2.A.1] in general). Theorem 3.11 (i) and Remark 3.12 (i) may be viewed as analogues of this classical statement for codimension 2 cycles.

(iii) To obtain a variant of Theorem 3.11 (iii) in the case where D is connected but not geometrically connected, one can apply Theorem 3.11 (iii) to the l -rational variety X_l over l , where l is the algebraic extension $l := H^0(D, \mathcal{O}_D)$ of k . The same remark applies to Theorem 3.11 (iv).

4. Smooth complete intersections of two quadrics

In this last section, we apply the above results to k -varieties X that are three-dimensional smooth complete intersections of two quadrics, computing the variety $\mathbf{CH}_{X/k}^2$ (in Theorem 4.5)

and providing a necessary and sufficient criterion for their k -rationality (in Theorem 4.7). A more classical necessary and sufficient criterion for their (separable) k -unirationality (Theorem 4.8) allows us to give examples, for any algebraically closed field κ , of such varieties over $\kappa((t))$ that are separably $\kappa((t))$ -unirational but not $\kappa((t))$ -rational (Theorem 4.14).

Many of the geometric results that we need along the way are available in the literature only in characteristic 0 or in characteristic $\neq 2$ [74, 27, 82], and we extend them to arbitrary characteristic.

4.1. Lines in a complete intersection of two quadrics

Let $X \subset \mathbf{P}_k^5$ be a three-dimensional smooth complete intersection of two quadrics over k . We let F be the Hilbert scheme of lines in X (also called the Fano variety of lines of X).

LEMMA 4.1. – *The following assertions hold:*

- (i) *The variety X does not contain any plane.*
- (ii) *The normal bundle $N_{L/X}$ of a line $L \subset X$ is isomorphic either to $\mathcal{O}_L^{\oplus 2}$ or to $\mathcal{O}_L(1) \oplus \mathcal{O}_L(-1)$.*
- (iii) *The variety F is a non-empty geometrically connected smooth projective surface with trivial canonical bundle. Its tangent space at a k -point corresponding to a line $L \subset X$ is naturally isomorphic to $H^0(X, N_{L/X})$.*

Proof of (i). – By the Lefschetz hyperplane theorem [42, XII, Corollary 3.7], $\text{Pic}(X)$ is generated by $\mathcal{O}_X(1)$. If X contained a plane P , we would have $\mathcal{O}_X(P) \simeq \mathcal{O}_X(l)$ for some $l \in \mathbf{Z}$, hence an equality of intersection numbers

$$1 = \mathcal{O}_P(1) \cdot \mathcal{O}_P(1) = \mathcal{O}_X(P) \cdot \mathcal{O}_X(1) \cdot \mathcal{O}_X(1) = 4l,$$

which is a contradiction.

Proof of (ii). – Since L and X are complete intersections in \mathbf{P}_k^5 , the normal exact sequence $0 \rightarrow N_{L/X} \rightarrow N_{L/\mathbf{P}_k^5} \rightarrow N_{X/\mathbf{P}_k^5}|_L \rightarrow 0$ of the inclusions $L \subset X \subset \mathbf{P}_k^5$ reads:

$$(4.1) \quad 0 \rightarrow N_{L/X} \rightarrow \mathcal{O}_L(1)^{\oplus 4} \rightarrow \mathcal{O}_L(2)^{\oplus 2} \rightarrow 0.$$

It follows that $N_{L/X}$ is a rank 2 vector bundle of degree 0 on L , hence is of the form $\mathcal{O}_L(l) \oplus \mathcal{O}_L(-l)$ for some $l \geq 0$ (see [47]). Since it admits an injective morphism to $\mathcal{O}_L(1)^{\oplus 4}$, one has $l \in \{0, 1\}$.

Proof of (iii). – The computation of the tangent space is [59, Chapter I, Theorem 2.8.1]. To prove that F is smooth, geometrically connected and non-empty, we may work over \bar{k} . For all $L \in F(\bar{k})$, one has $h^1(L, N_{L/X_{\bar{k}}}) = 0$ and $h^0(L, N_{L/X_{\bar{k}}}) = 2$ by (ii). The variety $F_{\bar{k}}$ is thus smooth of dimension 2 at L by [59, Chapter I, Theorem 2.8.3]. That $F_{\bar{k}}$ is non-empty and connected follows from [25, Théorème 2.1 b) c)]. To compute the canonical bundle of F , we let G be the Grassmannian of lines in $\mathbf{P}_{\bar{k}}^5$ and $0 \rightarrow \mathcal{S} \rightarrow \mathcal{O}_G^{\oplus 6} \rightarrow \mathcal{Q} \rightarrow 0$ be the short exact sequence of tautological bundles on G , where \mathcal{S} and \mathcal{Q} respectively have rank 4 and rank 2. The variety of lines F is defined in G by the vanishing of a section of $(\text{Sym}^2 \mathcal{Q})^{\oplus 2}$. Since F is smooth of the expected dimension, the normal short exact sequence reads

$0 \rightarrow T_F \rightarrow T_G|_F \rightarrow N_{F/G} \rightarrow 0$, where $N_{F/G} \simeq (\text{Sym}^2 \mathcal{Q})^{\oplus 2}|_F$ and $T_G|_F \simeq (\mathcal{S}^\vee \otimes \mathcal{Q})|_F$. We deduce that

$$K_F \simeq (\det((\text{Sym}^2 \mathcal{Q})^{\oplus 2}) \otimes \det(\mathcal{S} \otimes \mathcal{Q}^\vee))|_F \simeq \det(\mathcal{Q})|_F^{\otimes 6} \otimes \det(\mathcal{Q})|_F^{\otimes -6} \simeq \mathcal{O}_F. \quad \square$$

We will later show that $F_{\bar{k}}$ is actually an abelian surface (see Theorem 4.5).

Let $Z \subset X \times F$ be the universal line in X . If $\Lambda \subset X$ is a line, the second projection $Z \cap (\Lambda \times (F \setminus \{\Lambda\})) \rightarrow F \setminus \{\Lambda\}$ is a closed immersion by [36, Proposition 8.11.5]. Its image is a subscheme $W(\Lambda) \subseteq F \setminus \{\Lambda\}$ parametrizing the lines of X that are distinct from Λ and that intersect Λ . If $L \in W(\Lambda)(k)$ and $x = L \cap \Lambda$, then $T_L W(\Lambda) \subseteq T_L F = H^0(X, N_{L/X})$ is the subset of those $\sigma \in H^0(X, N_{L/X})$ such that $\sigma_x \in \langle T_x L, T_x \Lambda \rangle / T_x L$ (as can be seen using [76, Remark 4.5.4 (ii)]).

LEMMA 4.2. – *If L and Λ are two distinct lines in X that intersect, the inclusion of tangent spaces $T_L W(\Lambda) \subseteq T_L F$ is strict.*

Proof. – In view of Lemma 4.1 (ii), we may distinguish two cases according to the isomorphism class of the normal bundle $N_{L/X}$. Suppose first that $N_{L/X} \simeq \mathcal{O}_L^{\oplus 2}$. Consider the point $x = L \cap \Lambda$. Since $N_{L/X}$ is globally generated, we can choose a section $\sigma \in H^0(L, N_{L/X})$ such that $\sigma_x \notin \langle T_x L, T_x \Lambda \rangle / T_x L$. Then, the tangent vector of F at L associated to σ by Lemma 4.1 (iii) is not tangent to $W(\Lambda)$.

Assume now that $N_{L/X} \simeq \mathcal{O}_L(1) \oplus \mathcal{O}_L(-1)$. Choose homogeneous coordinates X_0, \dots, X_5 on \mathbf{P}_k^5 such that $L = \{X_0 = \dots = X_3 = 0\}$ and use X_0, \dots, X_3 to identify N_{L/\mathbf{P}_k^5} with $\mathcal{O}_L(1)^{\oplus 4}$. After a coordinate change, we may assume that the composition $\mathcal{O}_L(1) \rightarrow \mathcal{O}_L(1) \oplus \mathcal{O}_L(-1) \rightarrow \mathcal{O}_L(1)^{\oplus 4}$ of the inclusion of the first factor and of the first arrow of (4.1) is the inclusion of the first factor. In such coordinates,

$$\{X_0 + \varepsilon X_4 = X_1 = X_2 = X_3 = 0\} \text{ and } \{X_0 + \varepsilon X_5 = X_1 = X_2 = X_3 = 0\}$$

are two $k[\varepsilon]/(\varepsilon^2)$ -points of F . Assuming for contradiction that $T_L W(\Lambda) = T_L F$ and using the characterization recalled above of $T_L W(\Lambda)$ viewed as a subspace of $H^0(X, N_{L/X})$, we see that $\Lambda \subset P := \{X_1 = X_2 = X_3 = 0\}$. Since X contains L , the monomials $X_4^2, X_4 X_5$ and X_5^2 do not appear in the equations of X . Since X also contains the above two infinitesimal deformations of L , neither do the monomials $X_0 X_4$ and $X_0 X_5$. It follows that the intersection of X with the plane P is equal either to P or to the double line $\{X_0^2 = 0\} \subset P$. Since $\Lambda \subset X \cap P$ but $\Lambda \neq L$, we deduce that $P \subset X$, which contradicts Lemma 4.1 (i). \square

4.2. Projecting from a line

We keep the notation of §4.1.

Assume that X contains a line $\Lambda \subset X$, which we fix. We denote by $\mu : X' \rightarrow X$ and $\tilde{\mu} : (\mathbf{P}_k^5)' \rightarrow \mathbf{P}_k^5$ the blow-ups of Λ in X and in \mathbf{P}_k^5 , and by $\nu : X' \rightarrow \mathbf{P}_k^3$ and $\tilde{\nu} : (\mathbf{P}_k^5)' \rightarrow \mathbf{P}_k^3$ the morphisms obtained by projecting away from Λ .

PROPOSITION 4.3. – *There exists a smooth projective geometrically connected curve $\Delta \subset \mathbf{P}_k^3$ of genus 2 such that ν can be identified with the blow-up of Δ in \mathbf{P}_k^3 .*

Proof. – Write $\Lambda = \{X_0 = \dots = X_3 = 0\}$ in appropriate homogeneous coordinates X_0, \dots, X_5 of \mathbf{P}_k^5 . The morphism $\tilde{v} : (\mathbf{P}_k^5)' \rightarrow \mathbf{P}_k^3$ realizes $(\mathbf{P}_k^5)'$ as the projectivization (in Grothendieck’s sense) of the vector bundle $\mathcal{E} = \mathcal{O}_{\mathbf{P}_k^3} \oplus \mathcal{O}_{\mathbf{P}_k^3} \oplus \mathcal{O}_{\mathbf{P}_k^3}(1)$ on \mathbf{P}_k^3 . In a natural way, we use homogeneous coordinates X_0, \dots, X_3 on \mathbf{P}_k^3 , and we let X_4, X_5 (resp. X_6) denote the global sections of \mathcal{E} (resp. of $\mathcal{E}(-1)$) on \mathbf{P}_k^3 corresponding to the direct sum decomposition of \mathcal{E} . If X has equations $\{L_1X_4 + L_2X_5 + Q = 0\}$ and $\{L'_1X_4 + L'_2X_5 + Q' = 0\}$ in \mathbf{P}_k^5 , with L_1, L_2, L'_1, L'_2 (resp. Q, Q') linear (resp. quadratic) in X_0, \dots, X_3 , then X' has equations $\{L_1X_4 + L_2X_5 + QX_6 = 0\}$ and $\{L'_1X_4 + L'_2X_5 + Q'X_6 = 0\}$ in $(\mathbf{P}_k^5)'$.

The fibers of $v : X' \rightarrow \mathbf{P}_k^3$ are defined by two linear equations in a 2-dimensional projective space, hence are isomorphic to \mathbf{P}^0 , to \mathbf{P}^1 or to \mathbf{P}^2 . The determinantal subscheme $\Delta \subset \mathbf{P}_k^3$ defined by the vanishing of the maximal minors of the matrix

$$(4.2) \quad \begin{pmatrix} L_1 & L_2 & Q \\ L'_1 & L'_2 & Q' \end{pmatrix}$$

endows the subset of \mathbf{P}_k^3 over which the fibers of v are positive-dimensional with a schematic structure. We claim that $v|_{v^{-1}(\Delta)} : v^{-1}(\Delta) \rightarrow \Delta$ is a flat family of lines in the projective bundle $\tilde{v}|_{\tilde{v}^{-1}(\Delta)} : \tilde{v}^{-1}(\Delta) \rightarrow \Delta$. To see it, we work on an affine open subset of Δ with coordinate ring R . One has to show that the cokernel M of the linear map $R^2 \rightarrow R^3$ given by the transpose of the matrix (4.2) is free of rank 2. This follows from [28, Proposition 20.8] since $\text{Fitt}_1(M) = 0$ by the definition of Δ and $\text{Fitt}_2(M) = R$ by Lemma 4.1 (i).

Let us show that Δ is smooth of the expected dimension (equal to 1). To this end, we fix $x \in \Delta(\bar{k})$ such that $T_x\Delta_{\bar{k}}$ has dimension ≥ 2 and derive a contradiction.

We first assume that $v^{-1}(x)$ is the line $\{X_6 = 0\} \subset \tilde{v}^{-1}(x)$. Then the linear forms L_1, L_2, L'_1 and L'_2 vanish at x . As $T_x\Delta_{\bar{k}}$ has dimension ≥ 2 , the differentials at x of the cubic forms $L_1Q' - L'_1Q$ and $L_2Q' - L'_2Q$ are linearly dependent. After replacing (L_1, L'_1) and (L_2, L'_2) with suitable \bar{k} -linear combinations of (L_1, L'_1) and (L_2, L'_2) (which is possible by a change of coordinates), we may therefore assume that the cubic $\{L_1Q' - L'_1Q = 0\}$ is singular at x . Since Q and Q' do not both vanish at x (otherwise $X_{\bar{k}}$ would contain the plane $\tilde{\mu}(\tilde{v}^{-1}(x))$, contradicting Lemma 4.1 (i) over \bar{k}), and since L_1 and L'_1 vanish at x , we deduce that L_1 and L'_1 are linearly dependent. Consequently, an appropriate \bar{k} -linear combination of the degree 2 equations defining $X_{\bar{k}}$ is of the form $L''_2X_5 + Q''$, where L''_2 and Q'' are respectively linear and quadratic in X_0, \dots, X_3 . Since $V := \{L''_2X_5 + Q'' = 0\}$ is singular at $p := [0 : 0 : 0 : 0 : 1 : 0]$ and $X_{\bar{k}} \subset V$ is a Cartier divisor containing p , we see that $X_{\bar{k}}$ is also singular at p , which is absurd.

Thus the line $v^{-1}(x)$ is not equal to $\{X_6 = 0\} \subset \tilde{v}^{-1}(x)$. In other words, its image by μ is a line $L \subset X$ distinct from Λ . We note that the open subset of the relative Hilbert scheme of lines of $\tilde{v} : (\mathbf{P}_k^5)' \rightarrow \mathbf{P}_k^3$ consisting of those lines that are not defined by the equation $\{X_6 = 0\}$ in a fiber of \tilde{v} is naturally isomorphic to the scheme parametrizing the lines in \mathbf{P}_k^5 that are distinct from Λ but intersect Λ . Since moreover $v|_{v^{-1}(\Delta)} : v^{-1}(\Delta) \rightarrow \Delta$ is a family of lines over Δ , two independent tangent vectors of Δ at x give rise to two independent tangent vectors of $W(\Lambda)$ at L . As F is smooth of dimension 2 at L by Lemma 4.1 (iii), this contradicts Lemma 4.2 and finishes the proof that Δ is a smooth curve.

It then follows from [29, Theorem A2.60, Example A2.67] that \mathcal{O}_Δ is resolved by the Eagon-Northcott complex

$$0 \rightarrow \mathcal{O}_{\mathbf{P}_k^3}(-4)^{\oplus 2} \rightarrow \mathcal{O}_{\mathbf{P}_k^3}(-2) \oplus \mathcal{O}_{\mathbf{P}_k^3}(-3)^{\oplus 2} \rightarrow \mathcal{O}_{\mathbf{P}_k^3} \rightarrow \mathcal{O}_\Delta \rightarrow 0,$$

where the first arrow is given by the matrix (4.2) and the second one by its maximal minors. Using it allows one to compute that $h^0(\Delta, \mathcal{O}_\Delta) = 1$ and $h^1(\Delta, \mathcal{O}_\Delta) = 2$, hence that the smooth projective curve Δ is geometrically connected of genus 2.

To conclude, we denote by $\lambda : Y \rightarrow \mathbf{P}_k^3$ the blow-up of Δ in \mathbf{P}_k^3 .

Since $\nu|_{\nu^{-1}(\Delta)} : \nu^{-1}(\Delta) \rightarrow \Delta$ is a flat family of lines, hence a smooth morphism, the subscheme $\nu^{-1}(\Delta) \subset X'$ is a smooth divisor. Since X' is smooth, $\nu^{-1}(\Delta)$ is a Cartier divisor in X' , and the universal property of a blow-up yields a morphism $\phi : X' \rightarrow Y$ such that $\lambda \circ \phi = \nu$. Both λ and ν are birational (the latter because the generic fiber of ν is a 0-dimensional projective space), hence so is ϕ . We deduce that ϕ is an isomorphism, as is any birational morphism between smooth projective varieties with the same Picard number. \square

REMARK 4.4. – By Lemma 4.1 (iii), a smooth complete intersection of two quadrics in \mathbf{P}_k^5 contains a line over \bar{k} , hence is \bar{k} -rational by Proposition 4.3.

4.3. The intermediate Jacobian

In Theorem 4.5, we compute $\mathbf{CH}_{X/k}^2$ for threefolds X that are smooth complete intersections of two quadrics. Note that such varieties are \bar{k} -rational by Remark 4.4, so that Theorem 3.1 applies to them.

Theorem 4.5 (iv) will be used in a crucial manner in the proof of Theorem 4.7. In characteristic $\neq 2$, it goes back to the work of Wang [82].

In the statement of Theorem 4.5 (ii), we denote by $\text{Alb}_{V/k}^0$ (resp. $\text{Alb}_{V/k}^1$) the Albanese variety (resp. torsor) of a smooth proper geometrically connected variety V over k . This is the abelian variety over k (resp. torsor under $\text{Alb}_{V/k}^0$) which underlies the solution of the universal problem of morphisms from V to torsors under abelian varieties over k . We recall that $\text{Alb}_{V/k}^0$ is canonically dual to the abelian variety $(\mathbf{Pic}_{V/k}^0)^{\text{red}}$ and that the formation of $\text{Alb}_{V/k}^0$ is compatible with arbitrary extensions of scalars (see [33, Exp. 236, Théorème 3.3 (iii)], in which the geometric fibers of $X \rightarrow S$ should be assumed to be connected). We also recall that $\text{Alb}_{V/k}^0 = \mathbf{Pic}_{V/k}^0$ and $\text{Alb}_{V/k}^1 = \mathbf{Pic}_{V/k}^1$ if in addition V is a curve.

By a *conic* on X we mean a 1-dimensional closed subscheme of X which, when viewed as a subscheme of \mathbf{P}_k^5 , is the intersection of a quadric and a plane.

THEOREM 4.5. – *Let $X \subset \mathbf{P}_k^5$ be a smooth complete intersection of two quadrics, let F be its variety of lines, and let $Z \subset X \times F$ be the universal line.*

Denote by $\psi_X^2 : \text{CH}^2(X_{\bar{k}}) \xrightarrow{\sim} \mathbf{CH}_{X/k}^2(\bar{k})$ the isomorphism of Theorem 3.1 (iv). Then:

- (i) *The degree map $\text{deg} : \text{CH}^2(X_{\bar{k}}) \rightarrow \mathbf{Z}$ induces, via ψ_X^2 , a short exact sequence*

$$0 \rightarrow (\mathbf{CH}_{X/k}^2)^0 \rightarrow \mathbf{CH}_{X/k}^2 \xrightarrow{\delta} \mathbf{Z} \rightarrow 0$$
of k -group schemes.
- (ii) *The class $[\mathcal{O}_Z] \in \mathbf{K}_0(X_F)$ induces isomorphisms $F \xrightarrow{\sim} (\mathbf{CH}_{X/k}^2)^1 := \delta^{-1}(1)$ and $\text{Alb}_{F/k}^0 \xrightarrow{\sim} (\mathbf{CH}_{X/k}^2)^0$.*

- (iii) Let $\Xi \subset \mathbf{CH}^2(X_{\bar{k}})$ be the subset of classes represented by a conic on $X_{\bar{k}}$. There exists a unique reduced closed subscheme $D \subset (\mathbf{CH}_{X/k}^2)^2 := \delta^{-1}(2)$ such that $\psi_X^2(\Xi) = D(\bar{k})$. The scheme D is a smooth projective geometrically connected curve of genus 2 over k .
- (iv) Via the identifications $\mathbf{Pic}_{D/k}^0 = \text{Alb}_{D/k}^0$ and $\mathbf{Pic}_{D/k}^1 = \text{Alb}_{D/k}^1$, the inclusion $D \subset (\mathbf{CH}_{X/k}^2)^2$ induces isomorphisms of principally polarized abelian varieties $\mathbf{Pic}_{D/k}^0 \xrightarrow{\sim} (\mathbf{CH}_{X/k}^2)^0$ and of torsors $\mathbf{Pic}_{D/k}^1 \xrightarrow{\sim} (\mathbf{CH}_{X/k}^2)^2$.

Proof. – The sheaf \mathcal{O}_Z induces a class $[\mathcal{O}_Z] \in F_3\mathbf{G}_0(X_F) = F^2\mathbf{K}_0(X_F) = \mathbf{SK}_0(X_F)$ by §2.1.5 (as F is smooth by Lemma 4.1 (iii)). It therefore induces a morphism $F \rightarrow \mathbf{CH}_{X/k}^2$. This morphism factors through $(\mathbf{CH}_{X/k}^2)^1$ because it sends a point $x \in F(\bar{k})$ to the class in $\mathbf{CH}_{X/k}^2(\bar{k}) = \mathbf{CH}^2(X_{\bar{k}})$ of the line in $X_{\bar{k}}$ associated with x (as Z is flat over F), which has degree 1.

A morphism $a : F \rightarrow (\mathbf{CH}_{X/k}^2)^1$ having been constructed, we are now free to extend the scalars from k to any finite Galois extension of k : indeed, the existence, unicity and smoothness of D can be tested over such an extension, and all other conclusions of the theorem can even be tested over \bar{k} . As F is smooth, we may therefore, and will, assume that $F(k) \neq \emptyset$ (see [12, 2.2/13]).

Let us fix a line $\Lambda \subset X$ defined over k . Proposition 4.3 yields a diagram $X \xleftarrow{\mu} X' \xrightarrow{\nu} \mathbf{P}_k^3$, where μ is the blow-up of Λ and ν is the blow-up of a smooth projective geometrically connected curve $\Delta \subset \mathbf{P}_k^3$ of genus 2. Our knowledge of the Chow groups of a blow-up [31, Proposition 6.7 (e)] shows the existence of a Γ_k -equivariant short exact sequence $0 \rightarrow \mathbf{CH}^2(X_{\bar{k}})_{\text{alg}} \rightarrow \mathbf{CH}^2(X_{\bar{k}}) \xrightarrow{\text{deg}} \mathbf{Z} \rightarrow 0$. Assertion (i) now follows from (3.1) and (3.2).

Let $f : \text{Alb}_{F/k}^0 \rightarrow \text{Alb}_{(\mathbf{CH}_{X/k}^2)^1/k}^0 = (\mathbf{CH}_{X/k}^2)^0$ denote the morphism between Albanese varieties induced by $a : F \rightarrow (\mathbf{CH}_{X/k}^2)^1$. The morphism $b : \Delta \rightarrow F$ associating with $x \in \Delta$ the line $\mu(\nu^{-1}(x))$ induces a morphism $g : \mathbf{Pic}_{\Delta/k}^0 \rightarrow \text{Alb}_{F/k}^0$ between Albanese varieties.

The composition $f \circ g : \mathbf{Pic}_{\Delta/k}^0 \rightarrow (\mathbf{CH}_{X/k}^2)^0$ is an isomorphism of principally polarized abelian varieties. Indeed, it coincides at the level of \bar{k} -points with the principally polarized isomorphism $\mathbf{Pic}_{\Delta/k}^0 \xrightarrow{\sim} (\mathbf{CH}_{X/k}^2)^0$ obtained by applying Proposition 3.10 to μ and ν , hence is equal to it. It follows that the kernel of g is trivial and that $\text{Alb}_{F/k}^0$ has dimension ≥ 2 , hence that the first Betti number of $F_{\bar{k}}$ is ≥ 4 . Lemma 4.1 (iii) and the classification of surfaces with Kodaira dimension 0 (see [11, Table p.25 and Theorem 6]) now show that $F_{\bar{k}}$ is an abelian surface. We deduce that g is an isomorphism as its kernel is trivial and as $\mathbf{Pic}_{\Delta/k}^0$ and $\text{Alb}_{F/k}^0$ are both abelian surfaces. It follows that f is an isomorphism.

The variety F is isomorphic to an abelian variety since so is $F_{\bar{k}}$ and since $F(k) \neq \emptyset$. Thus a and f can be identified; hence a is an isomorphism as well, and (ii) is proved.

Let $c : \Delta \rightarrow (\mathbf{CH}_{X/k}^2)^2$ be defined by $c(x) = a(b(x)) + a(\Lambda)$, where Λ denotes the rational point of F corresponding to Λ . Passing to Albanese torsors yields a morphism $\mathbf{Pic}_{\Delta/k}^1 \rightarrow (\mathbf{CH}_{X/k}^2)^2$ which is an isomorphism since the underlying morphism of abelian varieties is the (principally polarized) isomorphism $f \circ g$. It follows, in particular, that c is a closed immersion.

Let us set $D = c(\Delta)$ and check that $D(\bar{k}) = \psi_X^2(\Xi)$, where $\Xi \subseteq \mathbf{CH}^2(X_{\bar{k}})$ is the subset appearing in (iii). The theorem will then be proved.

We take up the notation $\tilde{\mu}, \tilde{\nu}$ of §4.2 and the notation introduced in the proof of Proposition 4.3. The subvariety $\tilde{\mu}^{-1}(X) \subset (\mathbf{P}_k^5)'$ is given by the system of equations $\{(L_1X_4 + L_2X_5 + QX_6)X_6 = 0\}$ and $\{(L'_1X_4 + L'_2X_5 + Q'X_6)X_6 = 0\}$. If X_0, \dots, X_3 are the homogeneous coordinates of $x \in \Delta(\bar{k})$, this system of equations defines a conic in the plane $\tilde{\nu}^{-1}(x)$ since the matrix (4.2) has rank ≤ 1 . Thus, for any $x \in \Delta(\bar{k})$, we have exhibited a (singular) conic on $X_{\bar{k}}$ whose class in $\text{CH}^2(X_{\bar{k}})$ is visibly equal to $[\mu(\nu^{-1}(x))] + [\Lambda_{\bar{k}}]$, that is, to $(\psi_X^2)^{-1}(c(x))$. Hence $D(\bar{k}) \subseteq \psi_X^2(\Xi)$.

Let us now fix a conic C on $X_{\bar{k}}$ and prove that $\psi_X^2([C]) \in D(\bar{k})$. There exist a (unique) plane $P \subset \mathbf{P}_{\bar{k}}^5$ such that $C = X_{\bar{k}} \cap P$ and a (unique) quadric $Y \subset \mathbf{P}_{\bar{k}}^5$ containing $X_{\bar{k}} \cup P$. As X is smooth, the singular locus of Y is disjoint from $X_{\bar{k}}$ and has dimension ≤ 0 . Lemma 4.6 below provides a plane $P' \subset Y$ containing $\Lambda_{\bar{k}}$ such that $[P] = [P'] \in \text{CH}_2(Y)$. As $X_{\bar{k}}$ is an effective Cartier divisor on Y and as $X_{\bar{k}}$ contains neither P nor P' (see Lemma 4.1 (i)), it follows that $[C] = [X_{\bar{k}} \cap P] = [X_{\bar{k}} \cap P'] \in \text{CH}^2(X_{\bar{k}})$ [31, Proposition 2.6(a)]. Now $P' = \tilde{\mu}(\tilde{\nu}^{-1}(x))$ for a unique $x \in \mathbf{P}^3(\bar{k})$; as $X_{\bar{k}} \cap P'$ is a conic in P' , a glance at the equations of $\tilde{\mu}^{-1}(X) \cap \tilde{\nu}^{-1}(x)$ shows that (4.2) has rank ≤ 1 , hence $x \in \Delta(\bar{k})$ and $[X_{\bar{k}} \cap P'] = [\mu(\nu^{-1}(x))] + [\Lambda_{\bar{k}}]$, so that $\psi_X^2([C]) = c(x) \in D(\bar{k})$, as desired. \square

LEMMA 4.6. – *Let $Y \subset \mathbf{P}_{\bar{k}}^5$ be a quadric whose singular locus has dimension ≤ 0 . Let $P \subset Y$ be a plane. Let $\Lambda \subset Y$ be a line along which Y is smooth. There exists a (unique) plane $P' \subset Y$ rationally equivalent to P on Y such that $\Lambda \subset P'$.*

Proof. – Let X_0, \dots, X_5 denote the homogeneous coordinates of $\mathbf{P}_{\bar{k}}^5$. After a linear change of coordinates, we may assume that Λ is the line $X_0 = X_2 = X_4 = X_5 = 0$ and that Y is defined by the equation $X_0X_1 + X_2X_3 + X_4X_5 = 0$ (if Y is smooth) or by the equation $X_0X_1 + X_2X_3 + X_4^2 = 0$ (otherwise). Indeed, letting $H \subset \mathbf{P}_{\bar{k}}^5$ be a hyperplane containing Λ and avoiding the singular locus of Y , if Y is singular (so that Y is in this case a cone over the quadric $Y \cap H$, which is smooth), and setting $H = \mathbf{P}_{\bar{k}}^5$ otherwise, one can use Λ to split off two hyperbolic planes from a quadratic form defining the smooth quadric $Y \cap H$ [30, Proposition 7.13, Lemma 7.12]; the remaining regular quadratic form of dimension 1 or 2 has the desired shape since the ground field is algebraically closed.

If Y is smooth, the intersection of Y with the linear subspace $\{X_0 = X_2 = 0\}$ is the union of two planes containing Λ . According to *op. cit.*, Proposition 68.2, one of them is rationally equivalent to P on Y .

If Y is singular, let P' be the plane that contains Λ and the singular point of Y . The smooth quadric $Y \cap H$ cannot contain a plane (*op. cit.*, Lemma 8.10), therefore $P \cap H$ is a line. The two lines $P \cap H$ and Λ are rationally equivalent on $Y \cap H$ (*op. cit.*, Proposition 68.2); hence P and P' , being the cones over $P \cap H$ and over Λ inside Y , are rationally equivalent on Y [31, Example 2.6.2]. \square

4.4. Rationality

We can now prove a necessary and sufficient criterion for the k -rationality of three-dimensional smooth complete intersections of two quadrics.

THEOREM 4.7. – *Let $X \subset \mathbf{P}_k^5$ be a smooth complete intersection of two quadrics. Then X is k -rational if and only if it contains a line defined over k .*

Proof. – If X contains a line Λ , then projecting from Λ induces a birational map $X \dashrightarrow \mathbf{P}_k^3$ (see the more precise Proposition 4.3).

Assume, conversely, that X is k -rational. Let D be as in Theorem 4.5 (iii).

Let $\psi : (\mathbf{CH}_{X/k}^2)^0 \xrightarrow{\sim} \mathbf{Pic}_{D/k}^0$ be the inverse of the isomorphism of Theorem 4.5 (iv). Using Theorem 4.5 (ii), we identify the variety of lines F of X with the torsor $(\mathbf{CH}_{X/k}^2)^1$ under $(\mathbf{CH}_{X/k}^2)^0$.

Theorem 3.11 (iii) shows the existence of $d \in \mathbf{Z}$ such that $\psi_*[F] = [\mathbf{Pic}_{D/k}^d] \in H^1(k, \mathbf{Pic}_{D/k}^0)$ and Theorem 4.5 (iv) yields the identity $[\mathbf{Pic}_{D/k}^1] = 2\psi_*[F] \in H^1(k, \mathbf{Pic}_{D/k}^0)$. Combining these two equalities shows that

$$(4.3) \quad \psi_*[F] = [\mathbf{Pic}_{D/k}^d] = [\mathbf{Pic}_{D/k}^{1-d}] \in H^1(k, \mathbf{Pic}_{D/k}^0).$$

Noting that $K_D \in \mathbf{Pic}_{D/k}^2(k)$ since D has genus 2, we see that the $\mathbf{Pic}_{D/k}^0$ -torsor $\mathbf{Pic}_{D/k}^2$ is trivial. As one of d and $1-d$ is even, it follows from (4.3) that $\psi_*[F] = 0 \in H^1(k, \mathbf{Pic}_{D/k}^0)$. Consequently, $F(k) \neq \emptyset$ and X contains a line defined over k . \square

4.5. Unirationality

We now study the k -unirationality and the separable k -unirationality of smooth complete intersections of two quadrics.

THEOREM 4.8. – *Fix $N \geq 4$, and let $X \subset \mathbf{P}_k^N$ be a smooth complete intersection of two quadrics. The following assertions are equivalent:*

- (i) *The variety X is separably k -unirational.*
- (ii) *The variety X is k -unirational.*
- (iii) *One has $X(k) \neq \emptyset$.*

If $N = 4$, Theorem 4.8 (ii) \Leftrightarrow (iii) is due to Manin [67, Theorems 29.4 and 30.1] and Knecht [57, Theorem 2.1]. Over infinite perfect fields of characteristic not 2, Theorem 4.8 can be found in [17, Remark 3.28.3]. The counterpart of Theorem 4.8 (ii) \Leftrightarrow (iii) for cubic hypersurfaces is due to Kollár [60, Theorem 1.1], and we verify, in Theorem 4.13 below, that the equivalence with (i) holds for cubic hypersurfaces as well.

We start with a few lemmas. The first one is an analogue for separable k -unirationality of a statement proved by Kollár [60, Lemma 2.3] for k -unirationality.

LEMMA 4.9. – *Let X be an integral variety over k . The following are equivalent:*

- (i) *The variety X is separably k -unirational.*
- (ii) *The variety $X_{k(t)}$ is separably $k(t)$ -unirational.*
- (iii) *There exists a separable dominant rational map $\mathbf{A}_k^m \dashrightarrow X$ for some $m \geq 0$.*

Proof. – The implications (i) \Rightarrow (ii) \Rightarrow (iii) are immediate. To prove (iii) \Rightarrow (i), we adapt [60, Lemma 2.3]. We argue by induction on m . Let $\phi : \mathbf{A}_k^m \dashrightarrow X$ be a separable dominant rational map with $m > \dim(X)$, and let $U \subset \mathbf{A}_k^m$ be a dense open subset such that $\phi|_U$ is a smooth morphism. If k is infinite, set $k' = k$. If k is finite, choose a prime number ℓ invertible in k and a \mathbf{Z}_ℓ -extension k' of k .

As k' is infinite, one can choose a point $u = (u_1, \dots, u_m) \in U(k')$. After renumbering the u_i , we may assume that $k(u_m) \subseteq k(u_1)$, so that there exists $P_1 \in k[X_1]$ with $u_m = P_1(u_1)$. The ideal $I(u) \subset k[X_1, \dots, X_m]$ of u viewed as a closed point in \mathbf{A}_k^m is then generated by $X_m - P_1(X_1)$ and by elements of $k[X_1, \dots, X_{m-1}]$. It is thus generated by polynomials of the form $X_m - P(X_1, \dots, X_{m-1})$. Since u is étale over k , the zero locus $Z \subset \mathbf{A}_k^m$ of one of these polynomials intersects the fiber of $\phi|_U$ through u transversally at u . Then $\phi|_Z : Z \dashrightarrow X$ is smooth at u , hence it is dominant and separable. As $Z \simeq \mathbf{A}_k^{m-1}$, the induction hypothesis applies. \square

LEMMA 4.10. – *Let X be either a smooth complete intersection of two quadrics in \mathbf{P}_k^4 or a smooth cubic surface in \mathbf{P}_k^3 . In the cubic case, assume that the characteristic of k is not 2. Then there exists a hyperplane section of X that contains an ordinary double point.*

Proof. – Set $N = 4$ if X is an intersection of two quadrics in \mathbf{P}_k^4 and $N = 3$ if X is a cubic surface in \mathbf{P}_k^3 . By [67, Theorem 24.4], the variety X is isomorphic to a blow-up of $\mathbf{P}_{\bar{k}}^2$ in $9 - N$ points $p_1, \dots, p_{9-N} \in \mathbf{P}^2(\bar{k})$, no three of which lie on the same line. As the embedding $X \subset \mathbf{P}_k^N$ is given by the anticanonical bundle of X , the hyperplane sections of X correspond bijectively to the cubic curves in $\mathbf{P}_{\bar{k}}^2$ through the p_i . When $N = 4$, the hyperplane section associated with the union of the lines $(p_1 p_2)$ and $(p_3 p_4)$ and of a general line through p_5 has the required property. When $N = 5$ and the characteristic of k is not 2, the hyperplane section associated with the union of the unique conic through p_1, \dots, p_5 and of a general line through p_6 has the required property. (If k had characteristic 2, it could happen that a general line through p_6 is tangent to this conic.) \square

LEMMA 4.11. – *Fix $N \geq 4$, let $X \subset \mathbf{P}_k^N$ be a smooth complete intersection of two quadrics, and let $x \in X(\bar{k})$. Then, if $H \subset \mathbf{P}_k^N$ is a general hyperplane containing x , the variety $X \cap H$ is smooth of dimension $N - 3$.*

Proof. – Suppose for contradiction that the conclusion of the lemma does not hold. It follows that for every hyperplane $H \subset \mathbf{P}_k^N$ containing x , there exists a quadric $Q \subset \mathbf{P}_k^N$ in the pencil defining X such that $Q \cap H$ is not smooth of dimension $N - 2$ along some point of X . Consequently, the projective variety $W \subset \mathbf{P}(H^0(\mathbf{P}_k^N, \mathcal{O}(2))) \times \mathbf{P}(H^0(\mathbf{P}_k^N, \mathcal{O}(1)))$ parametrizing such pairs (Q, H) is at least $(N - 1)$ -dimensional.

Let $Q_0 \subset \mathbf{P}_k^N$ be any singular quadric in the pencil defining X . (Such a Q_0 exists since the locus of quadrics with a unique singular point has codimension 1 in the projective space of quadrics in \mathbf{P}_k^N , so that its closure contains an ample divisor.) Note that $x \in X \subset Q_0$. As X is smooth, the singular locus of Q_0 is zero-dimensional, hence Q_0 is a cone over a smooth quadric Q'_0 , and X does not contain the vertex of this cone. The projective dual $(Q_0)^\vee$ of Q_0 can be naturally identified with the projective dual of Q'_0 . By Lemma 4.12 below, the variety $(Q_0)^\vee$ has dimension $N - 2$ and is either a smooth quadric or a linear space whose dual is a line that meets Q_0 in its vertex and nowhere else. In both cases, the subset of $(Q_0)^\vee$ consisting of the hyperplanes that contain x is a non-trivial hyperplane section of $(Q_0)^\vee$, hence is $(N - 3)$ -dimensional. It follows that the variety parametrizing the hyperplanes $H \subset \mathbf{P}_k^N$ with $(Q_0, H) \in W$ is at most $(N - 3)$ -dimensional.

Let $\pi : W \rightarrow \mathbf{P}_k^1 \subset \mathbf{P}(H^0(\mathbf{P}_k^N, \mathcal{O}(2)))$ be the projection of W to the pencil of quadrics defining X . The fibers of π over singular quadrics have dimension $\leq N - 3$, and there exist such fibers. As $\dim(W) \geq N - 1$, some fiber of π over a smooth quadric must have dimension $\geq N - 1$. In other words, there exists a smooth quadric $Q \subset \mathbf{P}_k^N$ in the pencil defining X such that $(Q, H) \in W$ for all hyperplanes $H \subset \mathbf{P}_k^N$ containing x . Consequently, the projective dual of Q is the hyperplane dual to the point x . This contradicts Lemma 4.12 below since $x \in X \subset Q$. \square

LEMMA 4.12. – *Fix $N \geq 1$ and let $Q \subset \mathbf{P}_k^N$ be a smooth quadric. Then the projective dual of Q is a smooth quadric if k has characteristic $\neq 2$ or if N is odd; otherwise it is a hyperplane whose dual is a point not contained in Q .*

Proof. – In appropriate homogeneous coordinates, the quadric Q is defined by the equation $\{X_0X_1 + \cdots + X_{N-1}X_N = 0\}$ if N is odd, and by $\{X_0^2 + X_1X_2 + \cdots + X_{N-1}X_N = 0\}$ if N is even (see [40, XII, Proposition 1.2]). The lemma follows by a direct computation. \square

We finally reach the goal of this subsection.

Proof of Theorem 4.8. – That (i) implies (ii) is clear. Since projective k -unirational varieties always have k -points, (ii) implies (iii). It remains to show that (iii) implies (i). In view of Lemma 4.9, we may replace k with $k(t)$ to prove this implication. We thus assume from now on that k is infinite.

We argue by induction on $N \geq 4$. Suppose first that $N = 4$. Since k is infinite, [67, Theorems 29.4 and 30.1 (i)] shows that X is k -unirational. (The hypothesis that k is perfect is not used in the proof of [67, Theorem 30.1 (i)] for degree 4 del Pezzo surfaces. One only needs to notice that the variety of lines in a degree 4 del Pezzo surface over k is étale over k .) In particular, again because k is infinite, we see that $X(k)$ is Zariski dense in X . Choose $x \in X(k)$ general. Let $\nu : \tilde{X} \rightarrow X$ be the blow-up of X at x , with exceptional divisor $E \subset \tilde{X}$. By [67, Theorems 24.4 and 24.5] applied to $X_{\bar{k}}$ and $\tilde{X}_{\bar{k}}$, the linear system $|-K_{\tilde{X}}|$ yields an embedding of \tilde{X} in \mathbf{P}_k^3 as a cubic surface, the image of E being a line. Let $\pi : \tilde{X} \rightarrow \mathbf{P}_k^1$ be the conic bundle obtained by projecting \tilde{X} away from the line E . Choose $y \in \mathbf{P}^1(k)$ general. The scheme $\pi^{-1}(y) \cap E$ can be identified with the exceptional divisor of the blow-up at x of a hyperplane section of X that contains x and is singular at x . In view of Lemma 4.10, since x and y have been chosen general, the singularity at x of this hyperplane section is an ordinary double point (see [40, XVII, (4.1)]). We deduce that $\pi^{-1}(y) \cap E$ is smooth over k , hence that the morphism $\pi|_E : E \rightarrow \mathbf{P}_k^1$ is separable. The base change $Y := \tilde{X} \times_{\mathbf{P}_k^1} E \rightarrow E$ of π by $\pi|_E$ is a conic bundle over E with a section, hence is birational to $\mathbf{P}_k^1 \times E = \mathbf{P}_k^1 \times \mathbf{P}_k^1$. The projection $Y \rightarrow \tilde{X}$ is dominant and separable because so is $\pi|_E$. The variety \tilde{X} is thus separably k -unirational. This concludes the proof since X is birational to \tilde{X} .

If $N \geq 5$, choose $x \in X(k)$, and consider the space B of hyperplanes in \mathbf{P}_k^N that contain x . Let $Z = \{(w, b) \in X \times B; w \in b\}$ and let $p : Z \rightarrow B$ and $q : Z \rightarrow X$ be the natural projections. The generic fiber of p is a smooth complete intersection of two quadrics in $\mathbf{P}_{k(B)}^{N-1}$ by Lemma 4.11, and has a $k(B)$ -point induced by x , hence is separably $k(B)$ -unirational by the induction hypothesis. Since B is a projective space, it follows that Z is separably

k -unirational. As the generic fiber of the dominant map q is a projective space, hence is smooth, Lemma 4.9 shows that X is also separably k -unirational. \square

The strategy of the proof of Theorem 4.8 can also be applied to smooth cubic hypersurfaces, as we now briefly explain. Theorem 4.13 generalizes a theorem of Kollár, who proved the equivalence between (ii) and (iii) in [60, Theorem 1.1].

THEOREM 4.13. – *Fix $N \geq 3$, and let $X \subset \mathbf{P}_k^N$ be a smooth cubic hypersurface. The following assertions are equivalent:*

- (i) *The variety X is separably k -unirational.*
- (ii) *The variety X is k -unirational.*
- (iii) *One has $X(k) \neq \emptyset$.*

Proof. – The implication (i) \Rightarrow (ii) is trivial, and the equivalence (ii) \Leftrightarrow (iii) is [60, Theorem 1.1]. We prove (ii) \Rightarrow (i). By Lemma 4.9, we may assume that k is infinite. By (ii), one can thus choose $(x, y) \in (X \times X)(k)$ general. We argue by induction on $N \geq 3$. The induction step is identical to the one in the proof of Theorem 4.8, replacing Lemma 4.11 with Bertini’s theorem, which can be applied as x is general.

We now suppose that $N = 3$. Let C_x (resp. C_y) be the hyperplane section of X that is singular at x (resp. at y). By [60, First unirationality construction of §2, Proposition 3.1, Lemma 3.2], the curves C_x and C_y are k -rational, and the third intersection point map $\phi : C_x \times C_y \dashrightarrow X$ is dominant. As the map ϕ has degree 9, it is separable unless k has characteristic 3. Hence we may assume that k has characteristic $\neq 2$. Since (x, y) is general, one has $y \notin T_x X$, and the planes $T_x X$ and $T_y X$ intersect along a line D not containing y . If $x' \in C_x(k)$ is general, $T_{x'} C_x$ meets D at a unique point $z \neq y$. By Lemma 4.10 and [40, XVII, (4.1)], the curve C_y is nodal at y . One of the two tangent lines to C_y at y avoids z . We deduce that for $y' \in C_y(k)$ general, the line $T_{y'} C_y$ avoids z , hence does not meet $T_{x'} C_x$. This exactly means that ϕ is étale at (x', y') , hence separable. \square

4.6. Examples over fields of Laurent series

Here are examples of smooth complete intersections of two quadrics in $\mathbf{P}_{\kappa((t))}^5$ which are not $\kappa((t))$ -rational. The equations we use in characteristic 2 are borrowed from Bhosle [8].

THEOREM 4.14. – *Let κ be an algebraically closed field. If the characteristic of κ is $\neq 2$, let $a_0, \dots, a_5 \in \kappa$ be pairwise distinct elements, and consider the smooth projective variety $X \subset \mathbf{P}_{\kappa((t))}^5$ with equations*

$$\begin{cases} tX_0^2 + tX_1^2 + X_2^2 + \dots + X_5^2 = 0, \\ ta_0X_0^2 + ta_1X_1^2 + a_2X_2^2 + \dots + a_5X_5^2 = 0. \end{cases}$$

If κ has characteristic 2, let $a, b, c \in \kappa$ be pairwise distinct elements and consider the smooth projective variety $X \subset \mathbf{P}_{\kappa((t))}^5$ with equations

$$\begin{cases} tX_0X_1 + X_2X_3 + X_4X_5 = 0, \\ t(X_0^2 + aX_0X_1 + X_1^2) + (X_2^2 + bX_2X_3 + X_3^2) + (X_4^2 + cX_4X_5 + X_5^2) = 0. \end{cases}$$

Then X is separably $\kappa((t))$ -unirational, $\kappa((t^{\frac{1}{2}}))$ -rational, but not $\kappa((t))$ -rational.

Proof. – In view of Theorem 4.7 and Theorem 4.8, the conclusion of the theorem is equivalent to the assertion that X contains a point over $\kappa((t))$, a line over $\kappa((t^{\frac{1}{2}}))$, but no line over $\kappa((t))$, and this is what we shall now prove.

Let $Y \subset \mathbf{P}_\kappa^5$ be the subvariety with equations $\{\sum_i X_i^2 = \sum_i a_i X_i^2 = 0\}$ if κ has characteristic $\neq 2$, with equations

$$\begin{cases} X_0 X_1 + X_2 X_3 + X_4 X_5 = 0, \\ (X_0^2 + a X_0 X_1 + X_1^2) + (X_2^2 + b X_2 X_3 + X_3^2) + (X_4^2 + c X_4 X_5 + X_5^2) = 0 \end{cases}$$

if κ has characteristic 2. Let $\kappa((t)) \subset \kappa((u))$ be the quadratic extension with $u^2 = t$ and $\phi : X_{\kappa((u))} \xrightarrow{\sim} Y_{\kappa((u))}$ the isomorphism $(X_0, \dots, X_5) \mapsto (u X_0, u X_1, X_2, \dots, X_5)$.

As κ is algebraically closed, the variety X has a $\kappa((t))$ -point in the subspace $\{X_0 = X_1 = 0\}$. The variety Y contains a line (by Lemma 4.1 (iii)), so that $X_{\kappa((u))}$ contains a line as well. Let us now assume that X itself contains a line $L \subset X$, and derive a contradiction.

We denote by $\tau : \mu_2 \times \mathbf{P}_\kappa^5 \rightarrow \mathbf{P}_\kappa^5$ the action of the μ_2 -group scheme μ_2 on \mathbf{P}_κ^5 , via its non-trivial character on X_0 and X_1 and trivially on X_2, X_3, X_4 and X_5 . The subvariety $Y \subset \mathbf{P}_\kappa^5$ is τ -invariant. Let $\sigma : \mu_2 \times \text{Spec}(\kappa((u))) \rightarrow \text{Spec}(\kappa((u)))$ be the μ_2 -action endowing $\text{Spec}(\kappa((u)))$ with its natural structure of μ_2 -torsor over $\text{Spec}(\kappa((t)))$. It extends to an action of μ_2 on $\text{Spec}(\kappa[[u]])$ for which the closed point is an invariant subscheme.

Let us regard the line $L' := \phi(L_{\kappa((u))}) \subset Y_{\kappa((u))}$ as a $\kappa((u))$ -point of the variety of lines F of Y . In view of the equation defining ϕ and since L is defined over $\kappa((t))$, the morphisms $\mu_2 \times \text{Spec}(\kappa((u))) \rightarrow F$ given by the orbits of L' with respect to the actions of μ_2 on $F_{\kappa((u))}$ induced by τ and by σ coincide. It follows that the specialization $L'' \subset Y$ of L' with respect to the u -adic valuation of $\kappa((u))$ is τ -invariant. Since Y does not meet the projectivization of the subspace of κ^6 where μ_2 acts via its non-trivial character, the line L'' must be contained in the projectivization $\{X_0 = X_1 = 0\}$ of the subspace where μ_2 acts trivially. But the intersection of Y with $\{X_0 = X_1 = 0\}$ is an elliptic curve, which contains no line. This is the required contradiction, and the proposition is proved. \square

REMARKS 4.15. – (i) We do not know whether the variety X appearing in Proposition 4.14 is stably rational over $\kappa((t))$, even when $\kappa = \mathbf{C}$.

(ii) If κ has characteristic 2, the variety X considered in Proposition 4.14 is not $\kappa((t))$ -rational, but it becomes rational over the perfect closure $\kappa((t))_p$ of $\kappa((t))$. It follows that one cannot prove Proposition 4.14 by applying Theorem 4.7 over $\kappa((t))_p$. A theory of intermediate Jacobians over imperfect fields is therefore crucial for our proof of Theorem 4.14.

(iii) The variety X appearing in Proposition 4.14 when κ has characteristic 2 is the first example of a smooth projective variety over a field k which is k_p -rational, which has a k -point, but which is not k -rational. There are no such examples in dimension ≤ 2 (see Proposition 4.16 below).

(iv) Over fields k of characteristic $p > 2$, smooth complete intersections of two quadrics $X \subset \mathbf{P}_k^5$ cannot satisfy the conditions of Remark 4.15 (iii). Indeed, if X is k_p -rational, then the period of the variety of lines on X , viewed as a torsor under the intermediate Jacobian of X , must be a power of p , by Theorem 4.7 and [62, Proposition 5]; on the other hand, it divides 4 (see the proof of Theorem 4.7), hence it is equal to 1, so that X contains a line defined over k and is therefore k -rational. Adapting to imperfect fields the work of Kuznetsov

and Prokhorov [61] on prime Fano threefolds of genus 10 might lead to examples, over fields of characteristic 3, of smooth projective varieties that are k_p -rational, have a k -point, but are not k -rational. It remains a completely open problem whether varieties (resp. threefolds) satisfying these conditions exist in all characteristics $p \geq 5$.

(v) We do not know whether there exists a smooth projective variety over a separably closed field k which is \bar{k} -rational but not k -rational.

We include the following proposition, which generalizes [20, Theorem 1], to justify Remark 4.15 (iii).

PROPOSITION 4.16 (Segre, Manin, Iskovskikh). – *Let X be a smooth projective \bar{k} -rational surface over k . The following assertions are equivalent:*

- (i) X is k -rational,
- (ii) X is k_p -rational and $X(k) \neq \emptyset$.

If X is minimal, they are also equivalent to

- (iii) $K_X^2 \geq 5$ and $X(k) \neq \emptyset$.

Proof. – To prove the proposition, we may assume that X is minimal, from which it follows that X_{k_p} is minimal (see [73, Corollary 9.3.7]). That (i) implies (ii) is obvious. That (ii) and the minimality of X_{k_p} imply (iii) results from the birational classification of geometrically rational surfaces over perfect fields, due to Segre, Manin and Iskovskikh (see [50, p. 642]). It remains to prove that (iii) implies (i). If X is a del Pezzo surface, this is [81, Theorem 2.1]. Suppose now that X is not a del Pezzo surface. Then X belongs to the family II described in [49, Theorem 1]. Since $5 \leq K_X^2 \leq 8$ by [49, Theorem 3 (2)–(3)], one has $K_X^2 = 8$ by [49, Theorem 5]. It then follows from [49, Theorem 3 (2)] that X is either a product of curves of genus 0, or a projective bundle over a curve of genus 0. As $X(k) \neq \emptyset$, these curves of genus 0 are isomorphic to \mathbf{P}_k^1 , and X is k -rational. \square

BIBLIOGRAPHY

- [1] S. S. ABHYANKAR, *Resolution of singularities of embedded algebraic surfaces*, 2nd ed., Springer Monographs in Math., Springer, 1998.
- [2] D. ABRAMOVICH, K. KARU, K. MATSUKI, J. WŁODARCZYK, Torification and factorization of birational maps, *J. Amer. Math. Soc.* **15** (2002), 531–572.
- [3] J. D. ACHTER, S. CASALAINA-MARTIN, C. VIAL, On descending cohomology geometrically, *Compos. Math.* **153** (2017), 1446–1478.
- [4] J. ACHTER, S. CASALAINA-MARTIN, C. VIAL, A functorial approach to regular homomorphisms, preprint arXiv:1911.09911, to appear in *Algebraic Geometry*.

- [5] A. AUÉL, M. BERNARDARA, M. BOLOGNESI, Fibrations in complete intersections of quadrics, Clifford algebras, derived categories, and rationality problems, *J. Math. Pures Appl.* **102** (2014), 249–291.
- [6] A. BEAUVILLE, Variétés de Prym et jacobiniennes intermédiaires, *Ann. Sci. École Norm. Sup.* **10** (1977), 309–391.
- [7] O. BENOIST, O. WITTENBERG, The Clemens-Griffiths method over non-closed fields, *Algebr. Geom.* **7** (2020), 696–721.
- [8] U. BHOSLE, Pencils of quadrics and hyperelliptic curves in characteristic two, *J. reine angew. Math.* **407** (1990), 75–98.
- [9] S. BLOCH, Algebraic K -theory and algebraic geometry, in *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle Wash., 1972)*, Lecture Notes in Math. **341**, 1973, 259–265.
- [10] S. BLOCH, Torsion algebraic cycles and a theorem of Roitman, *Compos. Math.* **39** (1979), 107–127.
- [11] E. BOMBIERI, D. MUMFORD, Enriques’ classification of surfaces in char. p . II, in *Complex analysis and algebraic geometry*, 1977, 23–42.
- [12] S. BOSCH, W. LÜTKEBOHMERT, M. RAYNAUD, *Néron models*, *Ergeb. Math. Grenzgeb. (3)* **21**, Springer, 1990.
- [13] A. CHATZISTAMATIOU, K. RÜLLING, Vanishing of the higher direct images of the structure sheaf, *Compos. Math.* **151** (2015), 2131–2144.
- [14] C. CHEVALLEY, Une démonstration d’un théorème sur les groupes algébriques, *J. Math. Pures Appl.* **39** (1960), 307–317.
- [15] C. CLEMENS, P. GRIFFITHS, The intermediate Jacobian of the cubic threefold, *Ann. of Math.* **95** (1972), 281–356.
- [16] J.-L. COLLIOT-THÉLÈNE, J.-J. SANSUC, La descente sur les variétés rationnelles. II, *Duke Math. J.* **54** (1987), 375–492.
- [17] J.-L. COLLIOT-THÉLÈNE, J.-J. SANSUC, P. SWINNERTON-DYER, Intersections of two quadrics and Châtelet surfaces. I, *J. reine angew. Math.* **373** (1987), 37–107.
- [18] B. CONRAD, A modern proof of Chevalley’s theorem on algebraic groups, *J. Ramanujan Math. Soc.* **17** (2002), 1–18.
- [19] B. CONRAD, O. GABBER, G. PRASAD, *Pseudo-reductive groups*, second ed., New Mathematical Monographs **26**, Cambridge Univ. Press, 2015.
- [20] K. R. COOMBES, Every rational surface is separably split, *Comment. Math. Helv.* **63** (1988), 305–311.
- [21] V. COSSART, U. JANNSEN, S. SAITO, Canonical embedded and non-embedded resolution of singularities for excellent two-dimensional schemes, preprint arXiv:0905.2191.
- [22] V. COSSART, O. PILTANT, Resolution of singularities of threefolds in positive characteristic. I. Reduction to local uniformization on Artin-Schreier and purely inseparable coverings, *J. Algebra* **320** (2008), 1051–1082.

- [23] V. COSSART, O. PILTANT, Resolution of singularities of threefolds in positive characteristic. II, *J. Algebra* **321** (2009), 1836–1976.
- [24] O. DEBARRE, Polarisation sur les variétés abéliennes produits, *C. R. Acad. Sci. Paris Sér. I Math.* **323** (1996), 631–635.
- [25] O. DEBARRE, L. MANIVEL, Sur la variété des espaces linéaires contenus dans une intersection complète, *Math. Ann.* **312** (1998), 549–574.
- [26] M. DEMAZURE, P. GABRIEL, *Groupes algébriques. Tome I: Géométrie algébrique, généralités, groupes commutatifs*, Masson & Cie, Éditeur; North-Holland Publishing Co., 1970.
- [27] R. DONAGI, Group law on the intersection of two quadrics, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **7** (1980), 217–239.
- [28] D. EISENBUD, *Commutative algebra*, Graduate Texts in Math. **150**, Springer, 1995.
- [29] D. EISENBUD, *The geometry of syzygies*, Graduate Texts in Math. **229**, Springer, 2005.
- [30] R. ELMAN, N. KARPENKO, A. MERKURJEV, *The algebraic and geometric theory of quadratic forms*, American Mathematical Society Colloquium Publications **56**, Amer. Math. Soc., 2008.
- [31] W. FULTON, *Intersection theory*, second ed., *Ergeb. Math. Grenzgeb.* (3) **2**, Springer, 1998.
- [32] A. GROTHENDIECK, Éléments de géométrie algébrique: III. Étude cohomologique des faisceaux cohérents, I, *Publ. Math. IHES* **11** (1961).
- [33] A. GROTHENDIECK, *Fondements de la géométrie algébrique [extraits du Séminaire Bourbaki, 1957–1962]*, Secrétariat mathématique, 1962.
- [34] A. GROTHENDIECK, Éléments de géométrie algébrique: III. Étude cohomologique des faisceaux cohérents, II, *Publ. Math. IHES* **17** (1963).
- [35] A. GROTHENDIECK, Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas, II, *Publ. Math. IHES* **24** (1965).
- [36] A. GROTHENDIECK, Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas, III, *Publ. Math. IHES* **28** (1966).
- [37] A. GROTHENDIECK, *Schémas en groupes (SGA 3) II: Groupes de type multiplicatif, et structure des schémas en groupes généraux*, *Lecture Notes in Math.* **152**, Springer, 1970.
- [38] A. GROTHENDIECK, *Théorie des intersections et théorème de Riemann–Roch (SGA 6)*, *Lecture Notes in Math.* **225**, Springer, 1971.
- [39] A. GROTHENDIECK, *Théorie des topos et cohomologie étale des schémas (SGA 4 II)*, *Lecture Notes in Math.* **270**, Springer, 1972.
- [40] A. GROTHENDIECK, *Groupes de monodromie en géométrie algébrique (SGA 7 II)*, *Lecture Notes in Math.* **340**, Springer, 1973.
- [41] A. GROTHENDIECK, *Revêtements étales et groupe fondamental (SGA 1)*, *Documents Mathématiques* **3**, Soc. Math. France, 2003.

- [42] A. GROTHENDIECK, *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*, Documents Mathématiques 4, Société Mathématique de France, 2005.
- [43] A. GROTHENDIECK, *Schémas en groupes (SGA 3) I: Propriétés générales des schémas en groupes*, Documents Mathématiques 7, Société Mathématique de France, 2011.
- [44] R. HARTSHORNE, *Algebraic geometry*, Graduate Texts in Math. 52, Springer, 1977.
- [45] B. HASSETT, Y. TSCHINKEL, Cycle class maps and birational invariants, preprint arXiv:1908.00406, to appear in *Comm. Pure Appl. Math.*
- [46] B. HASSETT, Y. TSCHINKEL, Rationality of complete intersections of two quadrics, preprint arXiv:1903.08979, to appear in *L'Enseign. Math.*
- [47] M. HAZEWINKEL, C. F. MARTIN, A short elementary proof of Grothendieck's theorem on algebraic vector bundles over the projective line, *J. Pure Appl. Algebra* 25 (1982), 207–211.
- [48] L. ILLUSIE, M. TEMKIN, Exposé X: Gabber's modification theorem (log smooth case), *Astérisque* 363-364, 2014, 167–212.
- [49] V. A. ISKOVSKIKH, Minimal models of rational surfaces over arbitrary fields, *Izv. Akad. Nauk SSSR Ser. Mat.* 43 (1979), 19–43.
- [50] V. A. ISKOVSKIKH, Factorization of birational mappings of rational surfaces from the point of view of Mori theory, *Uspekhi Mat. Nauk* 51 (1996), 3–72.
- [51] U. JANNSEN, Weights in arithmetic geometry, *Japan J. Math.* 5 (2010), 73–102.
- [52] A. J. DE JONG, Smoothness, semi-stability and alterations, *Publ. Math. IHES* 83 (1996), 51–93.
- [53] J.-P. JOUANLOU, Riemann-Roch sans dénominateurs, *Invent. math.* 11 (1970), 15–26.
- [54] B. KAHN, On the universal regular homomorphism in codimension 2, preprint 2018, to appear in *Ann. Inst. Fourier*, <https://doi.org/10.5802/aif.3408>.
- [55] S. L. KLEIMAN, Geometry on Grassmannians and applications to splitting bundles and smoothing cycles, *Publ. Math. IHES* 36 (1969), 281–297.
- [56] S. L. KLEIMAN, The Picard scheme, in *Fundamental algebraic geometry*, Math. Surveys Monogr. 123, Amer. Math. Soc., 2005, 235–321.
- [57] A. KNECHT, Degree of unirationality for del Pezzo surfaces over finite fields, *J. Théor. Nombres Bordeaux* 27 (2015), 171–182.
- [58] F. F. KNUDSEN, D. MUMFORD, The projectivity of the moduli space of stable curves. I. Preliminaries on “det” and “Div”, *Math. Scand.* 39 (1976), 19–55.
- [59] J. KOLLÁR, *Rational curves on algebraic varieties*, *Ergeb. Math. Grenzgeb. (3)* 32, Springer, 1996.
- [60] J. KOLLÁR, Unirationality of cubic hypersurfaces, *J. Inst. Math. Jussieu* 1 (2002), 467–476.
- [61] A. KUZNETSOV, Y. PROKHOROV, Rationality of Fano threefolds over non-closed fields, *Amer. J. Math.* 145 (2023), 335–411.

- [62] S. LANG, J. TATE, Principal homogeneous spaces over abelian varieties, *Amer. J. Math.* **80** (1958), 659–684.
- [63] G. LAUMON, Homologie étale, in *Séminaire de géométrie analytique (École Norm. Sup., Paris, 1974-75)*, Astérisque **36–37**, 1976, 163–188.
- [64] J. LIPMAN, Notes on derived functors and Grothendieck duality, in *Foundations of Grothendieck duality for diagrams of schemes*, Lecture Notes in Math. **1960**, Springer, 2009, 1–259.
- [65] J. LIPMAN, A. NEEMAN, Quasi-perfect scheme-maps and boundedness of the twisted inverse image functor, *Illinois J. Math.* **51** (2007), 209–236.
- [66] Y. MANIN, Rational surfaces over perfect fields, *Publ. Math. IHES* **30** (1966), 55–113.
- [67] Y. MANIN, *Cubic forms*, second ed., North-Holland Mathematical Library **4**, North-Holland Publishing Co., 1986.
- [68] J. S. MILNE, *Étale cohomology*, Princeton Mathematical Series **33**, Princeton Univ. Press, N.J., 1980.
- [69] J. S. MILNE, Jacobian varieties, in *Arithmetic geometry (Storrs, Conn., 1984)*, Springer, 1986, 167–212.
- [70] J. P. MURRE, On contravariant functors from the category of pre-schemes over a field into the category of abelian groups (with an application to the Picard functor), *Publ. Math. IHES* **23** (1964), 5–43.
- [71] J. P. MURRE, Reduction of the proof of the non-rationality of a non-singular cubic threefold to a result of Mumford, *Compos. Math.* **27** (1973), 63–82.
- [72] J. P. MURRE, Applications of algebraic K -theory to the theory of algebraic cycles, in *Algebraic geometry, Sitges (Barcelona), 1983*, Lecture Notes in Math. **1124**, Springer, 1985, 216–261.
- [73] B. POONEN, *Rational points on varieties*, Graduate Studies in Math. **186**, Amer. Math. Soc., 2017.
- [74] M. REID, The complete intersection of two or more quadrics, Ph.D. Thesis, Cambridge University.
- [75] J. RIOU, Exposé XVI: Classes de Chern, morphismes de Gysin, pureté absolue, Astérisque **363-364**, 2014, 301–349.
- [76] E. SERNESI, *Deformations of algebraic schemes*, Grundlehren der math. Wiss. **334**, Springer, 2006.
- [77] J-P. SERRE, Appendix to Geometric methods for improving the upper bounds on the number of rational points on algebraic curves over finite fields, by K. Lauter, *J. Algebraic Geom.* **10** (2001), 19–36.
- [78] H. TANAKA, Behavior of canonical divisors under purely inseparable base changes, *J. reine angew. Math.* **744** (2018), 237–264.
- [79] R. W. THOMASON, Les K -groupes d'un schéma éclaté et une formule d'intersection excédentaire, *Invent. math.* **112** (1993), 195–215.

- [80] R. W. THOMASON, T. TROBAUGH, Higher algebraic K -theory of schemes and of derived categories, in *The Grothendieck Festschrift, Vol. III*, Progr. Math. **88**, Birkhäuser, 1990, 247–435.
- [81] A. VÁRILLY-ALVARADO, Arithmetic of del Pezzo surfaces, in *Birational geometry, rational curves, and arithmetic*, Simons Symp., Springer, 2013, 293–319.
- [82] X. WANG, Maximal linear spaces contained in the based loci of pencils of quadrics, *Algebr. Geom.* **5** (2018), 359–397.

(Manuscrit reçu le 16 décembre 2019 ;
accepté le 1^{er} août 2021.)

Olivier BENOIST
Département de mathématiques et applications
École normale supérieure et CNRS
45, rue d'Ulm
75230 Paris Cedex 05, France
E-mail: olivier.benoist@ens.fr

Olivier WITTENBERG
Institut Galilée
Université Sorbonne Paris Nord
99, avenue Jean-Baptiste Clément
93430 Villetaneuse, France
E-mail: wittenberg@math.univ-paris13.fr