


4 Properties of the CRT

Let \mathbf{A}_n be the set of all rooted oriented trees with n edges. If $\tau \in \mathbf{A}_n$ and $k \in \{0, \dots, 2n-1\}$ we define $\text{Reroot}(\tau, k)$ the tree τ re-rooted at the k -th corner in the counter clockwise contour of τ . See figure below.



Exercise 4.1. Let $k \in \{0, \dots, 2n-1\}$.

1. Show that $\text{Reroot}(\cdot, k) : \mathbf{A}_n \rightarrow \mathbf{A}_n$ is a bijection. Deduce that if τ_n is uniform over \mathbf{A}_n then $\text{Reroot}(\tau_n, k)$ is also uniform over \mathbf{A}_n .

Let $\tau \in \mathbf{A}_n$ and denote its contour function by $(C_n(t), t \in [0, 2n])$. We also write for $x \in [0, 1]$, $\mathbf{C}_n(x) = C_n(2nx)$. For every $r \geq 0$ let $\bar{r} = r - \lfloor r \rfloor$ be the fractional part of r . If $g : [0, 1] \rightarrow [0, +\infty[$ such that $g(0) = g(1) = 0$ and $s, t \in [0, 1]$ we recall the notation

$$m_g(s, t) = \inf \{g(u), u \in [s \wedge t, s \vee t]\}.$$

2. Show that the contour function $(\mathfrak{C}_n(x), x \in [0, 2n])$ of the rooted oriented tree $\text{Reroot}(\tau, k)$ is given for every $t \in [0, 1]$ by

$$\mathfrak{C}_n(t) = \mathbf{C}_n\left(\frac{k}{2n}\right) + \mathbf{C}_n\left(\frac{\overline{t+k}}{2n}\right) - 2m_{\mathbf{C}_n}\left(\frac{k}{2n}, \frac{\overline{k+t}}{2n}\right).$$

3. Deduce from the previous considerations that if $(\mathbf{e}(t))_{t \in [0, 1]}$ is a normalized Brownian excursion and $x \in [0, 1]$ then the process $(\mathfrak{e}(t))_{t \in [0, 1]}$ defined by

$$\mathfrak{e}(t) = \mathbf{e}(x) + \mathbf{e}(\overline{x+t}) - 2m_{\mathbf{e}}(x, \overline{x+t}),$$

has the same distribution as $(\mathbf{e}(t))_{t \in [0, 1]}$. What does it imply for the Brownian Continuum Random Tree ?


Let \mathcal{T} be a real tree. For $x \in \mathcal{T}$, the *multiplicity* of x is the number of connected components of $\mathcal{T} \setminus \{x\}$. A *leaf* is a point of multiplicity one. In the following $(\mathbf{e}(t))_{t \in [0, 1]}$ denotes a normalized Brownian excursion and $\mathcal{T}_{\mathbf{e}}$ its associated \mathbb{R} -tree, in particular we denote $p_{\mathbf{e}} : [0, 1] \rightarrow \mathcal{T}_{\mathbf{e}}$ the canonical projection.

Exercise 4.2. 1. Show that almost surely $p_{\mathbf{e}}(0)$ is a leaf of $\mathcal{T}_{\mathbf{e}}$. Deduce that for every $x \in [0, 1]$, $p_{\mathbf{e}}(x)$ is almost surely a leaf of $\mathcal{T}_{\mathbf{e}}$.

2. Show that a.s. the local minima of a Brownian motion are pairwise distinct.

3. Deduce that almost surely $\mathcal{T}_{\mathbf{e}}$ has only countable points of multiplicity 3 but no point of multiplicity strictly larger than 3.

Exercise 4.3. *Who are these charming gentlemen ?*



References

- [LG06] Jean-François Le Gall. Random real trees. *Ann. Fac. Sci. Toulouse Math. (6)*, 15(1):35–62, 2006.
- [MM06] Jean-François Marckert and Abdelkader Mokraddem. Limit of normalized quadrangulations: the Brownian map. *Ann. Probab.*, 34(6):2144–2202, 2006.